arXiv:1302.4049v1 [stat.AP] 17 Feb 2013

Angular Spectra for non-Gaussian Isotropic Fields

Gyorgy Terdik
Faculty of Informatics, University of Debrecen, Hungary,
Terdik.Gyorgy@inf.unideb.hu

February 19, 2013

Abstract

Cosmic Microwave Background (CMB) Anisotropies is a subject of intensive research in several fields of sciences [HD02]. In this paper we start a systematic development of basic notions and theory in statistics according to the application for CMB. The main result of this paper is the necessary and sufficient condition for isotropy of a non-Gaussian field in terms of spectra. Clear formulae between bi-, tri- and polyspectra and bi-, tri-, and higher order covariances are also given.

Keywords: Bispectrum, Trispectrum, Angular poly-Spectra, Cosmic microwave background radiation; Gaussianity; spherical random fields

1 Introduction

"Cosmological Principle (first coined by Einstein): the Universe is, in the large, homogeneous and isotropic" [Bar99]

In the last decade or so there has been some growing interest in studying the spatial - time data measured on the surface of a sphere. These data includes cosmic microwave background (CMB) anisotropies [KK06], [OH02], [AH12], medical imaging [KBID93] [Kak12], global and land-based temperature data [Jon94], [SRT06], gravitational and geomagnetic data, climate model [NC81].

One of the problem in focus is the Non-Gaussianity of the observations, which leads to the investigation of higher order angular spectra called polyspectra, [Hu01], [BSMR10], [BLSH12]. Angular polyspectra, in particular the bispectrum and trispectrum shows to be an appropriate measure of Gaussianity since for a Gaussian process all higher (then second order) spectra are zero, [KSH11]. An other important question is the Monte Carlo simulation of non-Gaussian isotropic maps with a given power spectrum and bispectrum and possible polyspectrum, [CM01], [RHS$^+$05].

In this paper some general properties of the angular polyspectra will be given for isotropic stochastic processes on sphere. Our treatment follows the basic theory of higher order spectra for non-Gaussian time series, [Bri65], [Bri01], [SRG84], [Ter99]. The symmetry relations and the appropriate series expansion of cumulants might have influence on the estimation of polyspectra as well, [SK12], [IMK12].

The isotropy assumption implies very particular form for the angular spectra therefore a delicate question is the construction of isotropic stochastic maps on the sphere.

2 Gaussian isotropic fields

Gaussian isotropic processes on sphere has a long history starting with [Obu47], some basic theory and references can be found in [Yag61], [Jon63], [McL86] [Yad83] [Yag87]. Due to the expansive recent applications there are new books [GG10], [CW11] and several papers covering a number of problems in general for spatial-time observations.

We consider a stochastic process $X(L)$ on the unit sphere $\mathbb{S}_2$ in $\mathbb{R}^3$, where $L = (\vartheta, \varphi)$, the co-latitude $\vartheta \in [0, \pi]$ and the longitude $\varphi \in [0, 2\pi]$. Let us suppose that $X(L)$ is continuous (in mean square sense), then it has a series expansion in terms of spherical harmonics Y_ℓ^m

$$X(L) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} Z_\ell^m Y_\ell^m(L), \tag{1}$$

where the coefficients are given by

$$Z_\ell^m = \int_{\mathbb{S}_2} X(L) \overline{Y_\ell^m(L)} \Omega(dL),$$

where $\Omega(dL) = \sin\vartheta d\vartheta d\varphi$ is Lebesque element of surface area on $\mathbb{S}_2$. It is generally accepted, in time series analysis that $X(L)$ called linear if Z_ℓ^m are independent and identically distributed ($EX(L) = 0$). Now it is straightforward that from the linearity here follows that the covariance function $\mathcal{C}(L_1, L_2) = \mathsf{E}X(L_1) X(L_2)$ depends on the angular distance γ between L_1 and L_2 only. Necessarily the covariance function depends on the central angle between locations and have the form

$$\mathcal{C}(L_1, L_2) = \mathcal{C}(\cos\gamma) = \sum_{\ell=0}^{\infty} f_\ell \frac{2\ell + 1}{4\pi} P_\ell(\cos\gamma), \tag{2}$$

where P_ℓ denotes the Legendre polynomial, see [Yad83], and all coefficients $f_\ell \geq 0$. Note here that the inner product $L_1 \cdot L_2 = \cos\gamma$. In other words $\mathcal{C}(L_1, L_2)$ is invariant under the group of rotations, i.e. $X(L)$ is *isotropic*. Moreover the assumption of linearity, i.e. the independence of the triangular array Z_ℓ^m so strong that the Gaussianity also follows. In other words the only linear field on the sphere is the Gaussian one. We shall consider a linearity which looks weaker although it will be shown to be equivalent to the whole independence of Z_ℓ^m.

Definition 1 *The field $X(L)$ is linear if the generating array Z_ℓ^m is uncorrelated and for fixed degree ℓ, $\{Z_\ell^m | m = -\ell, -\ell+1 \ldots, \ell-1, \ell\}$ are independent.*

This concept of linearity corresponds to the physical approach to the theory of angular momentum. One may consider linearity in some really weaker sense, namely the rows

$\{Z_\ell^m\}_{m=-\ell}^{\ell}$ of the array Z_ℓ^m, are independent but the variables Z_ℓ^m inside the row ℓ are uncorralated.

The convergence of the series$\sum_{\ell=0}^{\infty} f_\ell \frac{2\ell+1}{4\pi}$ is equivalent to the continuity of $\mathcal{C}(\cdot)$ on $[-1,1]$. The superposition (2) corresponds to the superposition of a covariance function on the real line in terms of its spectrum and the orthogonal system $\{\exp(i2\pi\lambda k), \quad k=0,1,2\ldots\}$, hence we treat ℓ as the frequency and $\widehat{\mathcal{C}}_\ell = f_\ell$ is the value of the spectrum at ℓ. Since $P_\ell(\cos 0) = 1$, the variance $\mathsf{E}X(L)^2$ is broken up into the sum of spectra therefore we have the analysis of variance. The inversion

$$\widehat{\mathcal{C}}_\ell = \int_{\mathbb{S}_2} \mathcal{C}(\cos\gamma) P_\ell(\cos\gamma)\,\Omega(dL),$$

is also valid. The stochastic process $X(L)$ itself has spectral representation (1) also in terms of spherical harmonics Y_ℓ^m with complex values, detailed account for spherical harmonics Y_ℓ^m is found in [VMK88], [SW71]. For a given $X(L)$ we have the inversion

$$Z_\ell^m = \int_{\mathbb{S}_2} X(L)\,\overline{Y_\ell^m(L)}\,\Omega(dL).$$

The orthogonal random 'measure' Z_ℓ^m is a triangular array for each fix ℓ; $m = -\ell, -\ell+1\ldots, \ell-1, \ell$, i.e. rows contain $2\ell+1$, i.i.d Gaussian random variables, $EZ_\ell^m = 0$, $EZ_\ell^m Z_k^{n*} = f_\ell \delta_{\ell,k}\delta_{m,n}$. The reason that f_ℓ does not depend on m, is the Funk-Hecke formula 23, see [Yad83], [Jon63].

The $\mathcal{C}(L_1 \cdot L_2)$ is strictly positive definite if all f_ℓ is ≥ 0, and only finitely many are zero ([Sch97], [Sch42]).

Example 2 *Matérn Class of Covariance Functions [Mat86], [SG05] Suppose one has a covariance function on the real line then its restriction to* $[0,\pi]$ *gives a covariance function* $\mathcal{C}_0(\gamma)$, *on the sphere* $\mathbb{S}_2$, $\mathcal{C}_0(\gamma) = \mathcal{C}(\cos\gamma)$, *for instance*

$$\begin{aligned}
\mathcal{C}_0(\gamma) &= \sigma^2 \frac{1}{\Gamma(\nu)} \left(\frac{\vartheta\gamma}{2}\right)^{\nu} 2K_\nu(\vartheta\gamma), \ \vartheta > 0, \ \nu > 0, \\
f_\ell &= \int_0^\pi \mathcal{C}_0(\gamma) \frac{2\ell+1}{4\pi} P_\ell(\cos\gamma)\,d\gamma.
\end{aligned}$$

Where K_ν *is the modified Bessel function of the second kind. Here the smoothness parameter* ν *controls the continuity, as well as* ϑ *controls the regularity [GG10], [She11], Note* $K_\nu(r) \sim \Gamma(\nu)(r/2)^{-\nu}/2$ *if* $r \to 0$, *and* $\nu > 0$, *hence* $\mathcal{C}_0(\gamma)/\sigma^2$ *is a correlation function.*

Example 3 *The generating function of Legendre polynomial* P_ℓ *is*

$$\sum_{\ell=0}^{\infty} P_\ell(x) z^\ell = \left(1 - 2xz + z^2\right)^{-1/2}, \ x \in (-1,1), \ |z| < 1$$

see 21, $0 < z < 1$, *hence*

$$\mathcal{C}(\cos\gamma) = \frac{1}{\sqrt{1 - 2z\cos\gamma + z^2}},$$

is covariance function on $[-1,1]$*, with* $f_\ell = \frac{4\pi}{2\ell+1} z^\ell$*. Similarly for* $n > 2$ *we have a covariance function*

$$\mathcal{C}(\cos\gamma) = \frac{1}{(1 - 2z\cos\gamma + z^2)^{(n-2)/2}},$$

for any $0 < z < 1$*.*

Example 4 *From the well known series*

$$\sum_{\ell=0}^{\infty} P_\ell(x) \frac{a^\ell}{2\ell+1} = \frac{1 - x^2}{(1 - 2xa + a^2)^{3/2}},\ x \in (-1,1),\ |a| < 1$$

see [Yad83], we find the covariance function on $[-1,1]$

$$\mathcal{C}(\cos\gamma) = \frac{1 - a^2}{(1 - 2a\cos\gamma + a^2)^{3/2}},$$

for any $0 < a < 1$*, and* $f_\ell = 4\pi a^\ell$*. Similarly for* $n \geq 2$*, we have*

$$\mathcal{C}(\cos\gamma) = \frac{1 - a^2}{(1 - 2a\cos\gamma + a^2)^{n/2}}.$$

Moreover the series

$$\sum_{\ell=1}^{\infty} P_\ell(x) \frac{t^\ell}{\ell} = \ln \frac{2}{1 - tx + \sqrt{1 - 2xt + t^2}},\quad |t| < 1,\ |x| < 1$$

see [PBM86], 5.10.1.4. Tom 2, gives a covariance function as well.

Example 5 *It is known, [Bre63], [WC01], that the probability density on the sphere*

$$\mathcal{C}(\cos\gamma) = \frac{\kappa}{4\pi\sinh(\kappa)} \exp(\kappa\cos\gamma),\ \kappa > 0$$

has series expansion

$$\mathcal{C}(\cos\gamma) = \frac{\kappa}{\sinh(\kappa)} \sum_{\ell=0}^{\infty} \frac{2\ell+1}{4\pi} \sqrt{\frac{\pi}{2\kappa}} I_{\ell+1/2}(\kappa) P_\ell(\cos\gamma),$$

where $\sqrt{\frac{\pi}{2\kappa}} I_{\ell+1/2}(\kappa)$ *is the modified spherical Bessel function of the first kind, [AS92], also*

$$(2\ell+1)\sqrt{\frac{\pi}{2\kappa}} I_{\ell+1/2}(\kappa) \sim \frac{\kappa^\ell}{(2\ell-1)!!}$$

see (3). Hence $\mathcal{C}$ *is a covariance function with spectrum*

$$\begin{aligned} f_\ell &= \frac{\kappa}{\sinh(\kappa)}\sqrt{\frac{\pi}{2\kappa}}I_{\ell+1/2}(\kappa) \\ &= \frac{I_{\ell+1/2}(\kappa)}{I_{1/2}(\kappa)}. \end{aligned}$$

Note $\sinh(\kappa)/\kappa = \sqrt{\frac{\pi}{2\kappa}}I_{1/2}(\kappa)$*, and*

$$\exp(a\cos\vartheta) = \sum_{\ell=0}^{\infty}(2\ell+1)\sqrt{\frac{\pi}{2a}}I_{\ell+1/2}(a)\,P_\ell(\cos\vartheta) \tag{3}$$

see [AS92], [PBM86].

Example 6 *For* $\kappa > 0$*,* $\gamma \in [0,\pi]$ *we have*

$$\sum_{\ell=0}^{\infty}\frac{\kappa^\ell}{\ell!}P_\ell(\cos\gamma) = \exp(\kappa\cos\gamma)\,J_0(\kappa\sin\gamma),$$

$$f_\ell = \frac{\kappa^\ell}{\ell!}\frac{4\pi}{2\ell+1},$$

see [PBM86], hence $\exp(\kappa\cos\gamma)\,J_0(\kappa\sin\gamma)/\exp(\kappa)$ *is a correlation function, where* J_0 *denotes Bessel function of the first kind.*

Example 7 *We have a covariance function of the form*

$$\begin{aligned} \mathcal{C}(\cos\gamma) &= I_0\left(2\kappa\sqrt{\cos\gamma-1}\right)I_0\left(2\kappa\sqrt{\cos\gamma+1}\right) \\ &= \sum_{\ell=0}^{\infty}\frac{\kappa^\ell}{(\ell!)^2}P_\ell(\cos\gamma), \\ f_\ell &= \frac{\kappa^\ell}{(\ell!)^2}\frac{4\pi}{2\ell+1} \end{aligned}$$

see [PBM86]. Note that $I_0(x)$ *is a function of* $x^2/4$ *see [AS92], hence there is no danger of complex values.* $I_0(0) = 1$*,* I_0 *denotes modified Bessel function of the first kind.*

Example 8 ***Poisson formula.*** *For a homogeneous isotropic field on* $\mathbb{R}^3$ *we have the spectral representation*

$$\mathcal{C}(r) = \int_0^\infty j_0(\lambda r)\,\Phi(d\lambda),$$

of a covariance function, [Yad83], and the restriction of $\mathcal{C}(r)$ *on the sphere* $\mathbb{S}_2$*,* $\mathcal{C}_0(\gamma) = \mathcal{C}(\cos\gamma)$*, is isotropic with spectrum*

$$f_\ell = 2\pi^2\int_0^\infty J^2_{\ell+1/2}(\lambda)\,\frac{1}{\lambda}\Phi(d\lambda). \tag{4}$$

Example 9 *For a homogeneous isotropic field on* $\mathbb{R}^{n+1}$, $n \geq 2$, *we have the spectral representation for a covariance function*

$$\mathcal{C}(r) = \int_0^\infty j_{n-3}(\lambda r)\, \Phi(d\lambda),$$

and the restriction of $\mathcal{C}(r)$ *on the sphere* $\mathbb{S}_n$, $\mathcal{C}(r) = \mathcal{C}_0(\cos\gamma)$, *is isotropic. The natural expansion of* $\mathcal{C}(r)$, *now is in terms of Gegenbauer polynomial* $\mathcal{C}_\ell^{(n-1)/2}(\cos\gamma)$ *[Yad83].*

Example 10 ***Laplace operator on*** $\mathbb{R}^3$. *Consider the homogeneous isotropic field on* $\mathbb{R}^3$ *according to the equation*

$$\left(\triangle - c^2\right)^m X = \partial W.$$

If $m = 1$, *its spectrum ([Yad83],* $n = 3$*), is*

$$S(\omega_1, \omega_2) = \frac{2}{(2\pi)^2} \frac{\omega^2}{\left(\omega^2 + c^2\right)^2}, \quad \omega^2 = \left\|(\omega_1, \omega_2)\right\|^2,$$

with covariance of Matérn Class

$$\begin{aligned}\mathcal{C}(r) &= \frac{1}{(2\pi)^{3/2}} \frac{K_{-1/2}(cr)}{2c\,(cr)^{-1/2}} \\ &= \frac{1}{(2\pi)^{3/2}} \frac{(cr)^{1/2} K_{1/2}(cr)}{2c},\end{aligned}$$

in terms of modified Bessel (Hankel) function, [AS92]. Now apply (4) for $\Phi(d\lambda) = S(\lambda)\, d\lambda$, *and we have*

$$\begin{aligned}f_\ell &= 2\pi^2 \int_0^\infty J_{\ell+1/2}^2(\lambda) \frac{1}{\lambda} \frac{2}{(2\pi)^2} \frac{\lambda^2}{\left(\lambda^2 + c^2\right)^2} d\lambda \\ &= \int_0^\infty J_{\ell+1/2}^2(\lambda) \frac{\lambda}{\left(\lambda^2 + c^2\right)^2} d\lambda,\end{aligned}$$

see [PBM86].

3 Non-Gaussian isotropy and the angular spectrum

In a physical phenomenon the isotropic property is treated as a principle. It means that there is no reason to make difference between directions. The corresponding property of a stochastic field is that the finite dimensional distributions remain unchanged after rotating the space.

3.1 Isotropy on sphere

Definition 11 *A stochastic field $X(L)$ on the unit sphere $\mathbb{S}_2$ is isotropic (in strict sense) if all finite dimensional distributions of $\{X(L), L \in \mathbb{S}_2\}$ are invariant under the rotation g for every $g \in SO(3)$.*

From now on we do not assume Gaussianity. But, for the simplicity, we suppose the existence of moments and that those determine the distribution as well. The general form of a mean square continuous field is given in terms of spherical harmonics

$$X(L) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} Z_\ell^m Y_\ell^m(L), \tag{5}$$

where $\{Z_\ell^m; \ell = 0, 1, \dots;\ m = -\ell, 1-\ell, \dots, -1, 0, 1, \dots, \ell-1, \ell\}$ is a triangular array, in the row ℓ we have $2\ell+1$ elements. Notice that $Y_0^0 = 1$, put $Z_0^0 = \mu$, otherwise $EZ_\ell^m = 0$, therefore $EX(L) = \mu$, and the convergence is valid in mean square sense [Leo99]. This development follows also from the stochastic Peter-Weyl Theorem, see [MP11] for details.

In case the m^{th} order cumulants of X are invariant under the rotation g for every $g \in SO(3)$ then it will be called *isotropic in m^{th}* order. Naturally a strictly isotropic field with m^{th} order moments is isotropic in m^{th} order.

We list here some properties which follows from isotropy, see [BM07] for details,

1. The distribution of both $\operatorname{Re} Z_\ell^m \overset{d}{=} \operatorname{Im} Z_\ell^m$ and $\operatorname{Re} Z_\ell^m / \operatorname{Im} Z_\ell^m$ are *Cauchy.*
2. Marginal distribution of both $\operatorname{Re} Z_\ell^m$ and $\operatorname{Im} Z_\ell^m$ are always symmetric
3. For fixed ℓ, Z_ℓ^m, $m = 0, 1, \dots, \ell-1, \ell$ are independent iff they are Gaussian.

Now, let us consider a rotation $g \in SO(3)$, it is known that the spherical harmonics Y_ℓ^m at the rotated location are given in terms of the Wigner D-matrix, see 18 Appendix B, more precisely

$$\Lambda(g) Y_\ell^m(L) = \sum_{k=-\ell}^{\ell} D_{k,m}^{(\ell)}(g) Y_\ell^k(L)$$

where $\Lambda(g)$ denotes the operator according to the rotation g, $\Lambda(g) Y_\ell^k(L) = Y_\ell^k\left(g^{-1}L\right)$. Hence the rotated field has the following form

$$\begin{aligned}\Lambda(g) X(L) &= \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} Z_\ell^m \sum_{k=-\ell}^{\ell} D_{k,m}^{(\ell)}(g) Y_\ell^k(L) \\ &= \sum_{\ell=0}^{\infty} \sum_{k=-\ell}^{\ell} \sum_{m=-\ell}^{\ell} D_{k,m}^{(\ell)}(g) Z_\ell^m Y_\ell^k(L).\end{aligned}$$

The isotropy assumption is equivalent to that the distribution of the variable

$$\mathcal{Z}_{\ell}^{k}\left(g\right)=\sum_{m=-\ell}^{\ell}D_{k,m}^{(\ell)}\left(g\right)Z_{\ell}^{m}, \tag{6}$$

is the same as the one of Z_{ℓ}^{k}. This statement will be used frequently below.

In matrix form

$$\mathcal{Z}_{\ell}\left(g\right)=D^{(\ell)}\left(g\right)Z_{\ell},$$

where $Z_{\ell}=\left[Z_{\ell}^{-\ell},Z_{\ell}^{-\ell+1},\ldots,Z_{\ell}^{\ell}\right]^{\top}$, $D^{(\ell)}=\left[D_{k,m}^{(\ell)}\right]_{-\ell,-\ell}^{\ell,\ell}$ (the dependence on g will be omitted unless it is necessary). This equation provides an equation for the cumulant functions Φ_{Z}^{ℓ} (log of the characteristic function) as well, in case of isotropy, for each rotation g we have

$$\Phi_{Z}^{\ell}\left(\underline{\omega}_{\ell}\right)=\Phi_{Z}^{\ell}\left(\underline{\omega}_{\ell}D^{(\ell)}\left(g\right)\right),$$

where $\underline{\omega}_{\ell}=\left[\omega_{-\ell},\omega_{-\ell+1},\ldots,\omega_{\ell-1},\omega_{\ell}\right]$. Hence the distribution of Z_{ℓ} should be rotationally invariant on $\mathbb{R}^{2\ell+1}$. For instance the covariance matrix $\mathcal{C}_{Z}\left(\ell_{1},\ell_{2}\right)=\operatorname{Cov}\left(Z_{\ell_{1}},Z_{\ell_{2}}\right)$ commutes with $D^{(\ell_{n})}$, since $D^{(\ell_{n})}$ is uniter and $\mathcal{C}_{Z}\left(\ell_{1},\ell_{2}\right)=D^{(\ell_{1})*}\mathcal{C}_{Z}\left(\ell_{1},\ell_{2}\right)D^{(\ell_{2})}$. If $\ell_{1}=\ell_{2}=\ell$, the only matrix which commutes with $D^{(\ell)}\left(g\right)$, for any $g\in SO\left(3\right)$ is constant times unit matrix.

Assume for a moment that $\operatorname{Cum}_{2}\left(Z_{\ell_{1}}^{k},Z_{\ell_{2}}^{m}\right)$ does depend on ℓ_{1},k,ℓ_{2},m, then we show below that from the isotropy $\operatorname{Cum}_{2}\left(\mathcal{Z}_{\ell_{1}}^{k},\mathcal{Z}_{\ell_{2}}^{m}\right)=\delta_{\ell_{1},\ell_{2}}\delta_{-k,m}\left(-1\right)^{m}\mathcal{C}_{2}\left(\ell_{1}\right)$ follows. Hence we should restrict ourselves on an uncorrelated generating array Z_{ℓ}^{m}, with $EZ_{\ell}^{m}Z_{k}^{n*}=\delta_{\ell,k}\delta_{m,n}\sigma_{\ell,m}^{2}$. Since $\sigma_{\ell,m}^{2}$ does not depend on m we denote it f_{ℓ}, moreover we shall consider the field X in the following form

$$X\left(L\right)=\sum_{\ell=0}^{\infty}\sum_{m=-\ell}^{\ell}Z_{\ell}^{m}Y_{\ell}^{m}\left(L\right),$$

where $EZ_{\ell}^{m}Z_{k}^{n*}=\delta_{\ell,k}\delta_{m,n}f_{\ell}$. In other words it is seen that the second and higher order structure of the generating process Z_{ℓ}^{m} inside the same degree ℓ are hiding, they are not identifiable.

According to the angular momentum of degree ℓ we define the field

$$u_{\ell}\left(L\right)=\sum_{m=-\ell}^{\ell}Z_{\ell}^{m}Y_{\ell}^{m}\left(L\right). \tag{7}$$

We shall be interested in the invariance of the distribution of $u_{\ell}\left(L\right)$ under rotations as well. If the location is fixed then $u_{\ell}\left(L\right)$ is connected to the distribution of $\mathcal{Z}_{\ell}^{k}$ directly, since the Condon and Shortley phase convention $\sqrt{\frac{2\ell+1}{4\pi}}D_{m,0}^{(\ell)}\left(g\right)=Y_{\ell}^{m*}\left(\vartheta,\varphi\right)$, where the rotation g given in Euler coordinates $\left(\gamma,\vartheta,\varphi\right)$, provides

$$u_{\ell}\left(L\right)=\sqrt{\frac{2\ell+1}{4\pi}}\sum_{m=-\ell}^{\ell}D_{0,m}^{(\ell)*}\left(\gamma,\vartheta,\varphi\right)Z_{\ell}^{m},$$

observe that γ here is arbitrary, see (32). Our main interest is the comparison of the finite dimensional distributions of the field $u_\ell(L)$ to the rotated one. In case a rotation carries the location L to the North pole N, u_ℓ simplifies

$$u_\ell(N) = \sqrt{\frac{2\ell+1}{4\pi}} Z_\ell^0.$$

Rewrite X into the form

$$X(L) = \sum_{\ell=0}^{\infty} u_\ell(L).$$

We consider a real valued $X(L)$, therefore $Z_\ell^{m*} \overset{d}{=} (-1)^m Z_\ell^{-m}$, since $Y_\ell^m(L)^* = (-1)^m Y_\ell^{-m}(L)$. Moreover if we reflect the location to the center then from $Y_\ell^m(-L) = (-1)^\ell Y_\ell^m(L)$ follows

$$X(-L) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} (-1)^\ell Z_\ell^m Y_\ell^m(L). \tag{8}$$

Therefore we have the following

Remark 12 *Since the parity $u_\ell(-L) = (-1)^\ell u_\ell(L)$ is valid, it implies that under assumption of isotropy*

$$\operatorname{Cum}_p\left(u_{\ell_1}(L_1), \ldots, u_{\ell_p}(L_p)\right) = (-1)^{\ell_1+\ell_2+\cdots+\ell_p} \operatorname{Cum}_p\left(u_{\ell_1}(L_1), \ldots, u_{\ell_p}(L_p)\right),$$

hence for $p > 2$ the sum $\mathcal{L}_p = \ell_1 + \ell_2 + \cdots + \ell_p$ must be even.

3.1.1 Second order Isotropy

Consider the second order cumulants of $\mathcal{Z}_\ell^k$, defined in (6),

$$\begin{aligned}
\operatorname{Cum}_2\left(\mathcal{Z}_{\ell_1}^k, \mathcal{Z}_{\ell_2}^m\right) &= \sum_{p,q=-\ell_1,\ell_2}^{\ell_1,\ell_2} D_{k,p}^{(\ell_1)} D_{m,q}^{(\ell_2)} \operatorname{Cum}_2\left(Z_{\ell_1}^p, Z_{\ell_2}^q\right) \\
&= \sum_{p,q=-\ell_1,\ell_2}^{\ell_1,\ell_2} D_{k,p}^{(\ell_1)} (-1)^p D_{m,q}^{(\ell_2)} \operatorname{Cum}_2\left(Z_{\ell_1}^{-p*}, Z_{\ell_2}^q\right).
\end{aligned}$$

Now assume isotropy, $\operatorname{Cum}_2\left(\mathcal{Z}_{\ell_1}^k, \mathcal{Z}_{\ell_2}^m\right) = \operatorname{Cum}_2\left(Z_{\ell_1}^k, Z_{\ell_2}^m\right)$ and integrate both sides over the sphere according to the invariant Haar measure, we have

$$\begin{aligned}
\operatorname{Cum}_2\left(Z_{\ell_1}^k, Z_{\ell_2}^m\right) &= \sum_{p,q=-\ell_1,\ell_2}^{\ell_1,\ell_2} \frac{\delta_{\ell_1,\ell_2}\delta_{-p,q}\delta_{-k,m}(-1)^k}{2\ell_1+1} \operatorname{Cum}_2\left(Z_{\ell_1}^{-p*}, Z_{\ell_2}^q\right) \\
&= \frac{\delta_{-k,m}(-1)^m}{2\ell_1+1} \sum_{p=-\ell_1}^{\ell_1} \delta_{\ell_1,\ell_2} \operatorname{Cum}_2\left(Z_{\ell_1}^{-p*}, Z_{\ell_2}^p\right) \\
&= \delta_{\ell_1,\ell_2}\delta_{-k,m}(-1)^m C_2(\ell_1)
\end{aligned}$$

where

$$C_2\left(\ell_1\right)=\frac{1}{2\ell_1+1}\sum_{p=-\ell_1}^{\ell_1}\operatorname{Cum}_2\left(Z_{\ell_1}^{-p*},Z_{\ell_1}^{p}\right).$$

Hence $\operatorname{Cum}_2\left(Z_{\ell_1}^{k},Z_{\ell_2}^{m*}\right)=\delta_{\ell_1,\ell_2}\delta_{k,m}f_\ell$, i.e. the series Z_ℓ^k is uncorrelated.

If the series Z_ℓ^k is uncorrelated then

$$\begin{aligned}\operatorname{Cum}_2\left(\mathcal{Z}_{\ell_1}^{k},\mathcal{Z}_{\ell_2}^{m}\right)&=\sum_{p,q=-\ell_1,\ell_2}^{\ell_1,\ell_2}D_{k,p}^{(\ell_1)}D_{m,q}^{(\ell_2)}\operatorname{Cum}_2\left(Z_{\ell_1}^{p},Z_{\ell_2}^{q}\right)\\&=\delta_{\ell_1,\ell_2}\sum_{p,q=-\ell_1}^{\ell_1}D_{k,p}^{(\ell_1)}D_{m,q}^{(\ell_1)}\left(-1\right)^p\operatorname{Cum}_2\left(Z_{\ell_1}^{-p*},Z_{\ell_1}^{q}\right)\\&=\delta_{\ell_1,\ell_2}\sum_{p=-\ell_1}^{\ell_1}D_{-p,-k}^{(\ell_1)*}D_{m,q}^{(\ell_1)}\left(-1\right)^k\delta_{-p,q}f_{\ell_1}\\&=\delta_{\ell_1,\ell_2}\delta_{-k,m}\left(-1\right)^m f_{\ell_1}\end{aligned}$$

since $D_{k,m}^{(\ell)}$ is unitary, hence $\operatorname{Cum}_2\left(\mathcal{Z}_{\ell_1}^{k},\mathcal{Z}_{\ell_2}^{-m*}\right)=\delta_{\ell_1,\ell_2}\delta_{-k,m}f_{\ell_1}=\operatorname{Cum}_2\left(Z_{\ell_1}^{k},Z_{\ell_2}^{-m*}\right)$.

Lemma 13 *The field $X\left(L\right)$ is isotropic in second order iff the triangular series Z_ℓ^k is uncorrelated with variance f_ℓ.*

We conclude that a field $X\left(L\right)$ with Gaussian i.i.d. Z_ℓ^m is strictly isotropic. It follows from the isotropy that u_{ℓ_1} and $u_{\ell_2}$with different degrees ℓ_1 and ℓ_2 are uncorrelated, and

$$\begin{aligned}\operatorname{Cum}_2\left(u_\ell\left(L_1\right),u_\ell\left(L_2\right)\right)&=\sum_{m_1,m_2=-\ell}^{\ell}\operatorname{Cum}_2\left(Z_\ell^{m_1},Z_\ell^{m_2}\right)Y_\ell^{m_1}\left(L_1\right)Y_\ell^{m_2}\left(L_2\right)\\&=\sum_{m_2=-\ell}^{\ell}f_\ell Y_\ell^{-m_2}\left(L_1\right)Y_\ell^{m_2}\left(L_2\right)\left(-1\right)^{m_2}\\&=f_\ell\sum_{m_2=-\ell}^{\ell}Y_\ell^{m_2*}\left(L_1\right)Y_\ell^{m_2}\left(L_2\right)\\&=\frac{2\ell+1}{4\pi}f_\ell P_\ell\left(L_1\cdot L_2\right),\end{aligned}$$

by the addition formula see (39). In particular $\operatorname{Var}\left(u_\ell\left(L\right)\right)=\frac{2\ell+1}{4\pi}f_\ell$. Instead of the addition formula one arrives to the same result if applies the isotropy for $\operatorname{Cum}_2\left(u_\ell\left(L_1\right),u_\ell\left(L_2\right)\right)$. Rotate the locations L_1 and L_2 such that L_2 becomes the North pole N and at the same time L_1 belongs to the plane xOz. This rotation will be denoted by $g_{L_2L_1}$. We have

$\mathrm{Cum}_2\left(u_\ell\left(L_1\right), u_\ell\left(L_2\right)\right) = \mathrm{Cum}_2\left(u_\ell\left(g_{L_2L_1}L_1\right), u_\ell\left(N\right)\right)$, since $u_\ell\left(N\right) = \sqrt{\frac{2\ell+1}{4\pi}}Z_\ell^0$ and $Y_\ell^0\left(\vartheta,\varphi\right) = \sqrt{\frac{2\ell+1}{4\pi}}P_\ell\left(\cos\vartheta\right)$,

$$\begin{aligned}
\mathrm{Cum}_2\left(u_\ell\left(L_1\right), u_\ell\left(L_2\right)\right) &= \mathrm{Cum}_2\left(u_\ell\left(g_{L_2L_1}L_1\right), \sqrt{\frac{2\ell+1}{4\pi}}Z_\ell^0\right)\\
&= \mathrm{Cum}_2\left(\sqrt{\frac{2\ell+1}{4\pi}}P_\ell\left(g_{L_2L_1}L_1\cdot N\right)Z_\ell^0, \sqrt{\frac{2\ell+1}{4\pi}}Z_\ell^0\right)\\
&= \frac{2\ell+1}{4\pi}f_\ell P_\ell\left(L_1\cdot L_2\right).
\end{aligned}$$

3.2 Spectrum

Consider the covariance function $\mathcal{C}\left(L_1, L_2\right) = \mathsf{E}\left(X\left(L_1\right)-\mu\right)\left(X\left(L_2\right)-\mu\right)$ for an isotropic field. Let the rotation $g_{L_2L_1}$ be the one which takes the location L_2 into the North pole N, and L_1 into the plane xOz. The Euler coordinates of $g_{L_2L_1}L_1$ is the co-latitude angle ϑ and 0,since the rotation does not change the angle ϑ between L_1 and L_2, such that $\cos\vartheta = L_1\cdot L_2$. Under isotropy assumption the joint distribution of $X\left(L_1\right)$ and $X\left(L_2\right)$ equals to the joint distribution of $X\left(N\right)$ and $X\left(g_{L_2L_1}L_1\right)$, i.e. $X\left(N\right)$ and $X\left(\vartheta, 0\right)$ contain all pairwise information. In other words the covariance of an isotropic field depends on ϑ only, $\mathrm{Cum}_2\left(X\left(L_1\right), X\left(L_2\right)\right) = \mathrm{Cov}\left(X\left(L_1\right), X\left(L_2\right)\right) = \mathcal{C}\left(g_{L_2L_1}L_1, N\right)$, necessarily $\mathcal{C}\left(L_1, L_2\right) = \mathcal{C}\left(L_1\cdot L_2\right) = \mathcal{C}\left(\cos\vartheta\right)$. Now

$$\begin{aligned}
X\left(N\right) &= \sum_{\ell=0}^{\infty}\sum_{m=-\ell}^{\ell} Y_\ell^m\left(N\right)Z_\ell^m\\
&= \sum_{\ell=0}^{\infty}\sum_{m=-\ell}^{\ell} \delta_{m,0}\sqrt{\frac{2\ell+1}{4\pi}}Z_\ell^m\\
&= \sum_{\ell=0}^{\infty}\sqrt{\frac{2\ell+1}{4\pi}}Z_\ell^0,
\end{aligned}$$

and

$$X\left(g_{L_2L_1}L_1\right) = \sum_{\ell=0}^{\infty}\sum_{m=-\ell}^{\ell} Y_\ell^m\left(g_{L_2L_1}L_1\right)Z_\ell^m,$$

We have $\mathrm{Cum}_2\left(Z_\ell^0, Z_k^n\right) = \delta_{\ell,k}\delta_{0,n}f_\ell$, hence

$$\begin{aligned}
\mathcal{C}\left(L_1, L_2\right) &= \sum_{\ell=0}^{\infty} f_\ell\sqrt{\frac{2\ell+1}{4\pi}}Y_\ell^0\left(g_{L_2L_1}L_1\right)\\
&= \sum_{\ell=0}^{\infty} f_\ell\frac{2\ell+1}{4\pi}P_\ell\left(\cos\vartheta\right).
\end{aligned}$$

Similarly to the time series setup the covariance $\mathcal{C}(L_1, L_2)$ is expanded in terms of an orthonormed system $\frac{2\ell+1}{4\pi}P_\ell(\cos\vartheta)$ with coefficients f_ℓ. In particular the variance $\operatorname{Var}(X(L))$ is split up into the superposition of f_ℓ's. Hence $S_2(\ell) = f_\ell$ is called **spectrum** of the field $X(L)$ according to the frequency ℓ.

4 Bispectrum

If the field $X(L)$ is non-Gaussian then the first characteristic after the second order moments to be considered is the third order cumulant, referred to bicovariance or 3-point covariance also since it is the third order central moment. The corresponding quantity in frequency domain is the bispectrum. Similarly to the spectrum when the covariance (second order cumulant)

$$\operatorname{Cum}_2(X(L_1), X(L_2)) = \sum_{\ell=0}^{\infty} f_\ell \frac{2\ell+1}{4\pi} P_\ell(\cos\vartheta),$$

is split up to the superposition of some orthogonal system of functions and the coefficients are called spectrum. We see that the orthogonal and normed system is the Legendre polynomial system $\left\{\frac{2\ell+1}{4\pi}P_\ell\right\}$ and the spectrum $S_2(\ell) = f_\ell$ corresponds to the frequency ℓ. We put a similar question for the higher order angular spectra as well. Namely, under assumption of isotropy of the m^{th} order cumulant we consider the series expansion according to an orthonormed system and the coefficients will be called m^{th} order spectrum. The Wigner 3j symbols

$$\begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & m_3 \end{pmatrix}$$

will be intensively used from now on, see Appendix B, 12. It depends on the quantum numbers ℓ_1, ℓ_2, ℓ_3 called degrees and orders m_1, m_2, m_3. We list some basic properties called *selection rules* for Wigner 3j symbols

- $3j$-symbol vanishes unless $m_1 + m_2 + m_3 = 0$,
- Integer perimeter rule $\ell_1 + \ell_2 + \ell_3$ is an integer and even if $m_1 = m_2 = m_3 = 0$,
- Triangular inequality $|\ell_1 - \ell_2| \le \ell_3 \le \ell_1 + \ell_2$ must be fulfilled.

4.1 Isotropy in third order

The rotation applied to he triangular array Z_ℓ^m shows the necessary and sufficient condition for the third order isotropy.

Lemma 14 *The field $X(L)$ is isotropic in third order iff the bicovariance $\operatorname{Cum}_3\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, Z_{\ell_3}^{m_3}\right)$ of the triangular series Z_ℓ^m has the form*

$$\operatorname{Cum}_3\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, Z_{\ell_3}^{m_3}\right) = \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & m_3 \end{pmatrix} B_3(\ell_1, \ell_2, \ell_3), \tag{9}$$

and

$$B_3(\ell_1,\ell_2,\ell_3) = \sum_{k_1,k_2,k_3} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ k_1 & k_2 & k_3 \end{pmatrix} \operatorname{Cum}_3\left(Z_{\ell_1}^{k_1}, Z_{\ell_2}^{k_2}, Z_{\ell_3}^{k_3}\right).$$

See Appendix C for the proof. Note here that $\operatorname{Cum}_3\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, Z_{\ell_3}^{m_3}\right) = 0$, if $m_1 + m_2 + m_3 \neq 0$, that is not all the possible bicovariances come into the picture, moreover the cumulants are depending on the orders m_1, m_2, m_3 through the Wigner 3j symbols only. In other words the third order probabilistic property inside a fixed quantum number ℓ does not show up. The function B_3 of the frequencies is an average of the cumulants $\operatorname{Cum}_3\left(Z_{\ell_1}^{k_1}, Z_{\ell_2}^{k_2}, Z_{\ell_3}^{k_3}\right)$ by 'probability' amplitudes, see (26) hence it is called as 'angle average bispectrum', [FS09], we shall turn back to the notion of the bispectrum later in this section.

Notice first that the quantity

$$\begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & m_3 \end{pmatrix} B_3(\ell_1,\ell_2,\ell_3),$$

is symmetric in ℓ_1, ℓ_2, ℓ_3, see Appendix C, C. More over $B_3(\ell_1,\ell_2,\ell_3)$ is symmetric in ℓ_1, ℓ_2, ℓ_3 as well, since by parity transformation (8) $\ell_1+\ell_2+\ell_3$ must be even, the coefficients $\begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ k_1 & k_2 & k_3 \end{pmatrix}$ are symmetric as well.

From now on we *fix the order* of ℓ_1, ℓ_2, ℓ_3 such that $\ell_1 \leq \ell_2 \leq \ell_3$.

Now we turn to the angular momentum field u_ℓ. First observe

$$\begin{aligned} \operatorname{Cum}_3\left(u_{\ell_1}(L_1), u_{\ell_2}(L_2), u_{\ell_3}(L_3)\right) &= \sum_{m_1,m_2,m_3} Y_{\ell_1}^{m_1}(L_1) Y_{\ell_2}^{m_2}(L_2) Y_{\ell_3}^{m_3}(L_3) \operatorname{Cum}_3\left(Z_{\ell_{1:3}}^{m_{1:3}}\right) \\ &= \sum_{m_1,m_2,m_3} Y_{\ell_1}^{m_1}(L_1) Y_{\ell_2}^{m_2}(L_2) Y_{\ell_3}^{m_3}(L_3) \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & m_3 \end{pmatrix} B_3(\ell_1,\ell_2,\ell_3) \\ &= B_3(\ell_1,\ell_2,\ell_3) \sum_{m_1,m_2,m_3} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & m_3 \end{pmatrix} Y_{\ell_1}^{m_1}(L_1) Y_{\ell_2}^{m_2}(L_2) Y_{\ell_3}^{m_3}(L_3). \end{aligned}$$

This is invariant both the rotation and ordering of ℓ_1, ℓ_2, ℓ_3. The *3-product of spherical harmonics* Y_ℓ^m,

$$\widetilde{I}_{\ell_1,\ell_2,\ell_3}(L_1,L_2,L_3) = \sum_{m_1,m_2,m_3} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & m_3 \end{pmatrix} Y_{\ell_1}^{m_1}(L_1) Y_{\ell_2}^{m_2}(L_2) Y_{\ell_3}^{m_3}(L_3)$$

is *rotational invariant*, see Appendix (30), therefore without any loss of generality we apply the rotation $g_{L_3 L_2}$ which takes the location L_3 into the North pole N and at the same time takes L_2 into the zOx- plane

$$\operatorname{Cum}_3\left(u_{\ell_1}\left(L_1\right), u_{\ell_2}\left(L_2\right), u_{\ell_3}\left(L_3\right)\right) = \operatorname{Cum}_3\left(u_{\ell_1}\left(g_{L_3L_2}L_1\right), u_{\ell_2}\left(g_{L_3L_2}L_2\right), u_{\ell_3}\left(N\right)\right)$$
$$= \sqrt{\frac{2\ell_3+1}{4\pi}} B_3\left(\ell_1,\ell_2,\ell_3\right) \sum_{m_1=-\ell_1}^{\ell_1} \sum_{m_2=-\ell_2}^{\ell_2} Y_{\ell_1}^{m_1}\left(g_{L_3L_2}L_1\right) Y_{\ell_2}^{m_2}\left(g_{L_3L_2}L_2\right) Y_{\ell_3}^{0}\left(N\right) \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & 0 \end{pmatrix}$$
$$= \sqrt{\frac{2\ell_3+1}{4\pi}} B_3\left(\ell_1,\ell_2,\ell_3\right) \sum_{m_1,m_2} Y_{\ell_1}^{m_1}\left(g_{L_3L_2}L_1\right) Y_{\ell_2}^{m_2}\left(g_{L_3L_2}L_2\right) \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & 0 \end{pmatrix}.$$

The third order cumulants of u_ℓ contains an orthonormed system of functions

$$I_{\ell_1,\ell_2,\ell_3}\left(L_1,L_2,L_3\right) = I_{\ell_1,\ell_2,\ell_3}\left(g_{L_3L_2}L_1, g_{L_3L_2}L_2, N\right)$$
$$= \sqrt{\frac{2\ell_3+1}{4\pi}} \sum_{m_{1:2}} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & 0 \end{pmatrix} Y_{\ell_1}^{m_1}\left(g_{L_3L_2}L_1\right) Y_{\ell_2}^{m_2}\left(g_{L_3L_2}L_2\right).$$

In this way $I_{\ell_1,\ell_2,\ell_3}\left(L_1,L_2,L_3\right)$ is connected to the *bipolar spherical harmonics*, i.e. to the irreducible tensor products of the spherical harmonics with different arguments, [VMK88].

$$\left[Y_{\ell_1}\left(L_1\right) \otimes Y_{\ell_2}\left(L_2\right)\right]_{\ell,m} = (-1)^{\ell_1-\ell_2+m} \sqrt{2\ell+1} \sum_{m_1,m_2} \begin{pmatrix} \ell_1 & \ell_2 & \ell \\ m_1 & m_2 & m \end{pmatrix} Y_{\ell_1}^{m_1}\left(L_1\right) Y_{\ell_2}^{m_2}\left(L_2\right)$$

Rewrite I_{ℓ_1,ℓ_2,ℓ_3} in terms of Euler angles

$$\mathcal{I}_{\ell_1,\ell_2,\ell_3}\left(\vartheta_1,\varphi_1,\vartheta_2\right) = I_{\ell_1,\ell_2,\ell_3}\left(g_{L_3L_2}L_1, g_{L_3L_2}L_2, N\right)$$
$$= \sqrt{\frac{2\ell_3+1}{4\pi}} \sum_{m_1,m_2} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & 0 \end{pmatrix} Y_{\ell_1}^{m_1}\left(\vartheta_1,\varphi_1\right) Y_{\ell_2}^{-m_1}\left(\vartheta_2,0\right)$$
$$= \sqrt{\frac{2\ell_3+1}{4\pi}} \sum_{m_1,m_2} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & 0 \end{pmatrix} (-1)^{m_1} Y_{\ell_1}^{m_1}\left(\vartheta_1,\varphi_1\right) Y_{\ell_2}^{m_1*}\left(\vartheta_2,0\right).$$

According to the usual measure $\Omega\left(dL_1\right)\Omega\left(dL_2\right) = \sin\vartheta_1 d\vartheta_1 d\varphi_1 \sin\vartheta_2 d\vartheta_2 d\varphi_2$ on $\vartheta \in [0,\pi]$, $\varphi \in [0,2\pi]$, we have

$$\int_{\mathbb{S}_2} Y_\ell^{m*}\left(\vartheta,0\right) Y_j^k\left(\vartheta,0\right) \Omega\left(dL\right) = \delta_{\ell,j}\delta_{m,k},$$

hence

$$\iint\limits_{\mathbb{S}_2} \mathcal{I}_{\ell_1,\ell_2,\ell_3}^{*} \mathcal{I}_{j_1,j_2,j_3} \Omega\left(dL_1\right)\Omega\left(dL_2\right) = \delta_{\ell_1,j_1}\delta_{\ell_2,j_2}\delta_{\ell_3,j_3},$$

since

$$\left(2\ell_3+1\right) \sum_{m_1,m_2} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & 0 \end{pmatrix}^2 = 1.$$

The third order cumulants of u_ℓ simplifies

$$\operatorname{Cum}_3\left(u_{\ell_1}\left(L_1\right), u_{\ell_2}\left(L_2\right), u_{\ell_3}\left(L_3\right)\right)=B_3\left(\ell_1, \ell_2, \ell_3\right) \mathcal{I}_{\ell_1, \ell_2, \ell_3}\left(\vartheta_1, \varphi_1, \vartheta_2\right).$$

If we denote it by

$$\operatorname{Cum}_3\left(u_{\ell_1}\left(L_1\right), u_{\ell_2}\left(L_2\right), u_{\ell_3}\left(L_3\right)\right)=C_{u, \ell_1, \ell_2, \ell_3}\left(\vartheta_1, \varphi_1, \vartheta_2\right),$$

then we have the inversion formula

$$\begin{aligned}&\iint_{\mathbb{S}_2} C_{u, j_1, j_2, j_3}\left(\vartheta_1, \varphi_1, \vartheta_2\right) \mathcal{I}_{\ell_1, \ell_2, \ell_3}\left(\vartheta_1, \varphi_1, \vartheta_2\right) \Omega\left(d L_1\right) \Omega\left(d L_2\right) \\ &=\delta_{\ell_1, j_1} \delta_{\ell_2, j_2} \delta_{\ell_3, j_3} B_3\left(\ell_1, \ell_2, \ell_3\right).\end{aligned}$$

4.2 Bispectrum of isotropic fields on sphere

The use of the bispectrum for detecting non-Gaussianity and nonlinearity is well known in time series analysis [SRG84], [Hin82] [TM98], the similar question has been put and studied for CMB analysis see [Mar04], [AH12] and references therein. The asymptotics of the bispectrum has a long history started by [RvN65], the asymptotic distribution of angular bispectrum when the degrees are increasing is considered by [Mar06], [Mar08].

Start with the general case $\operatorname{Cum}_3\left(X\left(L_1\right), X\left(L_2\right), X\left(L_3\right)\right)$. The value of the bicovariance does not change under the rotation $g_{L_3 L_2}$

$$\begin{aligned}\operatorname{Cum}_3\left(X\left(L_1\right), X\left(L_2\right), X\left(L_3\right)\right) &=\operatorname{Cum}_3\left(X\left(g_{L_3 L_2} L_1\right), X\left(g_{L_3 L_2} L_2\right), X(N)\right) \\ &=\operatorname{Cum}_3\left(X\left(\vartheta_1, \varphi_1\right), X\left(\vartheta_2, 0\right), X(N)\right),\end{aligned}$$

the spherical coordinates of the locations $g_{L_3 L_2} L_1=\left(\vartheta_1, \varphi_1\right)$, $g_{L_3 L_2} L_2=\left(\vartheta_2, 0\right)$ are defined by the locations L_1, L_2, L_3 as follows: $g_{L_3 L_2} L_2 \cdot N=L_2 \cdot L_3=\cos \vartheta_2$, $g_{L_3 L_2} L_1 \cdot N=L_1 \cdot L_3=\cos \vartheta_1$. These angles $\left(\vartheta_1, \varphi_1, \vartheta_2\right)$ define uniquely, up to rotations, the triangle given by locations L_1, L_2, L_3. In general the bicovariance is written $C_3\left(\vartheta_1, \varphi_1, \vartheta_2\right)=\operatorname{Cum}_3\left(X\left(\vartheta_1, \varphi_1\right), X\left(\vartheta_2, 0\right), X(N)\right)$, i.e. it depends on a location $L\left(\vartheta_2, 0\right)$ form the main circle $\left(\varphi_2=0\right)$ and a general location $L_0=L_0\left(\vartheta_1, \varphi_1\right)$.

$$\begin{aligned}&\operatorname{Cum}_3\left(X\left(L_1\right), X\left(L_2\right), X\left(L_3\right)\right) \\ &=\sum_{\ell_1, \ell_2, \ell_3=0}^{\infty} \sqrt{\frac{2 \ell_3+1}{4 \pi}} \sum_{m_1=-\ell_1}^{\ell_1} \sum_{m_2=-\ell_2}^{\ell_2} \operatorname{Cum}_3\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, Z_{\ell_3}^0\right) Y_{\ell_1}^{m_1}\left(g_{L_3 L_2} L_1\right) Y_{\ell_2}^{m_2}\left(g_{L_3 L_2} L_2\right) \\ &=\sum_{\ell_1, \ell_2, \ell_3=0}^{\infty} B_3\left(\ell_1, \ell_2, \ell_3\right) \mathcal{I}_{\ell_1, \ell_2, \ell_3}\left(\vartheta_1, \varphi_1, \vartheta_2\right).\end{aligned}$$

This series expansion of $\operatorname{Cum}_3\left(X\left(L_1\right), X\left(L_2\right), X\left(L_3\right)\right)$ implies the following definition

Definition 15 *The bispectrum of the isotropic field $X(L)$ is*

$$S_3(\ell_1,\ell_2,\ell_3) = B_3(\ell_1,\ell_2,\ell_3),$$

and the bicoherence of $X(L)$ is

$$\frac{S_3(\ell_1,\ell_2,\ell_3)}{\sqrt{S_2(\ell_1)\,S_2(\ell_2)\,S_2(\ell_3)}} = \frac{B_3(\ell_1,\ell_2,\ell_3)}{\sqrt{f_{\ell_1} f_{\ell_2} f_{\ell_3}}}.$$

4.2.1 Particular cases

For any locations L_1 and L we have

$$\mathrm{Cum}_3\left(u_{\ell_1}(L_1), u_{\ell_2}(L), u_{\ell_3}(L)\right) = \sqrt{\prod_i \frac{2\ell_i+1}{4\pi}} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ 0 & 0 & 0 \end{pmatrix} B_3(\ell_1,\ell_2,\ell_3)\, P_{\ell_1}(\cos L\cdot L_1),$$

from this readily follows

$$\begin{aligned}\mathrm{Cum}_3\left(X(L_1), X(L), X(L)\right) = \sum_{\ell_1,\ell_2,\ell_3=0}^{\infty} \prod_{i=1}^{3} \sqrt{\frac{2\ell_i+1}{4\pi}} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ 0 & 0 & 0 \end{pmatrix} \\ \times B_3(\ell_1,\ell_2,\ell_3)\, P_{\ell_1}(\cos L\cdot L_1).\end{aligned}$$

In particular, according to the Condon and Shortley phase convention, we have for any location L the third order central moment

$$\mathrm{Cum}_3\left(u_{\ell_1}(L), u_{\ell_2}(L), u_{\ell_3}(L)\right) = \sqrt{\prod \frac{2\ell_i+1}{4\pi}} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ 0 & 0 & 0 \end{pmatrix} B_3(\ell_1,\ell_2,\ell_3),$$

and the third order cumulant (central moment) $\mathrm{Cum}_3\left(X(L), X(L), X(L)\right)$ of $X(L)$ is

$$\begin{aligned}\mathrm{Cum}_3\left(X(L), X(L), X(L)\right) &= \mathrm{Cum}_3\left(X(N), X(N), X(N)\right) \\ &= \sum_{\ell_1,\ell_2,\ell_3=0}^{\infty} \mathrm{Cum}_3\left(Z_{\ell_1}^0, Z_{\ell_2}^0, Z_{\ell_3}^0\right) \prod \sqrt{\frac{2\ell_i+1}{4\pi}} \\ &= \sum_{\ell_1,\ell_2,\ell_3=0}^{\infty} \prod \sqrt{\frac{2\ell_i+1}{4\pi}} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ 0 & 0 & 0 \end{pmatrix} B_3(\ell_1,\ell_2,\ell_3)\end{aligned}$$

Note that the summation is taken over those ℓ_1, ℓ_2, ℓ_3 when $\ell_1+\ell_2+\ell_3 = \mathcal{L}$ is even and for this case the following explicit expression is valid

$$\begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ 0 & 0 & 0 \end{pmatrix} = (-1)^{\mathcal{L}/2} \sqrt{\frac{\prod(\mathcal{L}-2\ell_j)!}{(\mathcal{L}+1)!}} \frac{(\mathcal{L}/2)!}{\prod(\mathcal{L}/2-\ell_j)!}$$

4.3 Linear field

First we assume that the rows of the triangle array Z_ℓ^m contain independent variables. In other words the angular momentum field u_ℓ is linear. Then for a fixed degree ℓ

$$\mathrm{Cum}_3\left(Z_\ell^{m_1}, Z_\ell^{m_2}, Z_\ell^{m_3}\right) = \delta_{m_i=m}\begin{pmatrix}\ell & \ell & \ell\\ m & m & m\end{pmatrix} B_3\left(\ell,\ell,\ell\right),$$

now the selection rules apply, see Appendix B, 13

$$\mathrm{Cum}_3\left(Z_\ell^{m_1}, Z_\ell^{m_2}, Z_\ell^{m_3}\right) = \delta_{m_i=m}\delta_{m=0}\begin{pmatrix}\ell & \ell & \ell\\ 0 & 0 & 0\end{pmatrix} B_3\left(\ell,\ell,\ell\right).$$

Hence the only nonzero third order cumulant might be $\mathrm{Cum}_3\left(Z_\ell^0, Z_\ell^0, Z_\ell^0\right)$. Further the bispectrum

$$\begin{aligned} B_3\left(\ell,\ell,\ell\right) &= \sum_k \begin{pmatrix}\ell & \ell & \ell\\ k & k & k\end{pmatrix} \mathrm{Cum}_3\left(Z_\ell^k, Z_\ell^k, Z_\ell^k\right)\\ &= \begin{pmatrix}\ell & \ell & \ell\\ 0 & 0 & 0\end{pmatrix} \mathrm{Cum}_3\left(Z_\ell^0, Z_\ell^0, Z_\ell^0\right),\end{aligned}$$

therefore

$$\mathrm{Cum}_3\left(Z_\ell^0, Z_\ell^0, Z_\ell^0\right) = \begin{pmatrix}\ell & \ell & \ell\\ 0 & 0 & 0\end{pmatrix}^2 \mathrm{Cum}_3\left(Z_\ell^0, Z_\ell^0, Z_\ell^0\right).$$

We conclude from this that from the isotropy and independence follows $\mathrm{Cum}_3\left(Z_\ell^m, Z_\ell^m, Z_\ell^m\right) = 0$. Moreover if all the members of the triangle array Z_ℓ^m are independent then $\mathrm{Cum}_3\left(X\left(L_1\right), X\left(L_2\right), X\left(L_3\right)\right) = 0$. Third order cumulants are vanishes for instance when the distribution is Gaussian, or symmetric, etc.

4.4 Symmetries of the bispectrum

The cumulants $\mathrm{Cum}_3\left(X\left(L_1\right), X\left(L_2\right), X\left(L_3\right)\right)$ according to different locations L_1, L_2 and L_3 are defined by the spherical triangle with vertices L_1, L_2 and L_3. The efficiency of computations and avoiding the redundancy in statistical procedures require to determine the principal domains. The *principal domain* for the bispectrum according to the frequencies ℓ_1, ℓ_2, ℓ_3 is the following

1. ℓ_1, ℓ_2, ℓ_3 is monotone: $\ell_1 \leq \ell_2 \leq \ell_3$,
2. $\ell_1 + \ell_2 + \ell_3$ is even, see Remark 12,
3. ℓ_1, ℓ_2, ℓ_3 fulfils the triangular inequality $|\ell_1 - \ell_2| \leq \ell_3 \leq \ell_1 + \ell_2$,
4. $\mathrm{Cum}_3\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, Z_{\ell_3}^{m_3}\right) = 0$, unless $m_1 + m_2 + m_3 = 0$.

Similarly the *principal domain* for the bicovariance according to the locations (L_1, L_2, L_3) is $\{(\vartheta_1, \varphi_1, \vartheta_2)\,|\vartheta_1 \in [0,\pi], \varphi_2 \in [0,\pi], \vartheta_2 \in [0,\pi]\}$. We apply the following notation of these angles $\cos\vartheta_1 = L_2 \cdot L_3$, $\cos\vartheta_2 = L_1 \cdot L_3$, and $\varphi_2 = \phi_3$ is the surface angle at L_3. The third central angle is given by $\cos\vartheta_3 = L_1 \cdot L_2$. The surface angle ϕ_3 can be calculated for instance from the cosine formula

$$\cos\vartheta_3 = \cos\vartheta_1 \cos\vartheta_2 + \sin\vartheta_1 \sin\vartheta_2 \cos\varphi_2.$$

5 Trispectrum

5.1 Isotropy in fourth order

In the papers [Hu01], [KK06] for instance, the notion of *trispectrum* is used for the coefficients of the series expansion of the fourth order moments. Naturally, because of the expression of the moment via cumulants (18) it contains terms which correspond to the Gaussian part of the field. This is one of the reasons that the time series analysis considers the series expansion of cumulants instead of moments and reserve the notion of *trispectrum* for this case only, see [Bri65], [MH95]. The cumulants up to third order equal to the central moments, but it is not so for higher order, see Appendix A. Cumulants are very useful, for instance the measuring the distance of a distribution from the Gaussian one.

Lemma 16 *The field $X(L)$ is isotropic in fourth order iff the cumulant* $\operatorname{Cum}_4\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, Z_{\ell_3}^{m_3}, Z_{\ell_4}^{m_4}\right)$ *of the triangular array Z_ℓ^m has the form*

$$\begin{aligned}\operatorname{Cum}_4\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, Z_{\ell_3}^{m_3}, Z_{\ell_4}^{m_4}\right) = \sum_{\ell^1, m^1} (-1)^{m^1} \begin{pmatrix} \ell_1 & \ell_2 & \ell^1 \\ m_1 & m_2 & m^1 \end{pmatrix} \begin{pmatrix} \ell^1 & \ell_3 & \ell_4 \\ -m^1 & m_3 & m_4 \end{pmatrix} \\ \times \sqrt{2\ell^1+1}\, T_4\left(\ell_1, \ell_2, \ell_3, \ell_4 |\, \ell^1\right) \qquad (10)\end{aligned}$$

See the Appendix D for the proof. The function $T_4\left(\ell_1, \ell_2, \ell_3, \ell_4 |\, \ell^1\right)$ has the form

$$\begin{aligned}T_4\left(\ell_1, \ell_2, \ell_3, \ell_4 |\, \ell^1\right) = \sqrt{2\ell^1+1} \sum_{k_1,k_2,k_3,k_4,k^1} (-1)^{k^1} \begin{pmatrix} \ell_1 & \ell_2 & \ell^1 \\ k_1 & k_2 & k^1 \end{pmatrix} \begin{pmatrix} \ell^1 & \ell_3 & \ell_4 \\ -k^1 & k_3 & k_4 \end{pmatrix} \\ \times \operatorname{Cum}_4\left(Z_{\ell_1}^{k_1}, Z_{\ell_2}^{k_2}, Z_{\ell_3}^{k_3}, Z_{\ell_4}^{k_4}\right) \qquad (11)\end{aligned}$$

We notice here, the (11) shows that the cumulants of Z_ℓ^m and the function $T_4\left(\ell_1, \ell_2, \ell_3, \ell_4 |\, \ell^1\right)$ define each other uniquely.

Remark 17 *If we assume that $T_4\left(\ell_1, \ell_2, \ell_3, \ell_4 |\, \ell^1\right)$ is known for $\ell_1 \le \ell_2 \le \ell_3 \le \ell_4$, then the cumulants* $\operatorname{Cum}_4\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, Z_{\ell_3}^{m_3}, Z_{\ell_4}^{m_4}\right)$ *are given for any $(\ell_1, \ell_2, \ell_3, \ell_4)$.*
Vice versa $T_4\left(\ell_1, \ell_2, \ell_3, \ell_4 |\, \ell^1\right)$ can be calculated for any $(\ell_1, \ell_2, \ell_3, \ell_4)$ by (11).

The sum in (11) goes for k_1, k_2, k_3, k_4 and $k^1 = k_1 + k_2 = -k_3 - k_4$. Hence the summation contains those k_m's when equation $k_1 + k_2 + k_3 + k_4 = 0$, fulfils. Similarly $\mathrm{Cum}_4\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, Z_{\ell_3}^{m_3}, Z_{\ell_4}^{m_4}\right) = 0$, if $m_1 + m_2 + m_3 + m_4 \neq 0$. The triangular inequality for ℓ_1, ℓ_2, ℓ^1 and ℓ_3, ℓ_4, ℓ^1 suggests that the 'quadrilateral' with edges $(\ell_1, \ell_2, \ell_3, \ell_4)$ consists of two triangles ℓ_1, ℓ_2, ℓ^1 and ℓ_3, ℓ_4, ℓ^1. The cumulants are invariant of the order of their variables therefore this should be true for the other diagonal as well ℓ_4, ℓ_1, ℓ and ℓ_2, ℓ_3, ℓ. Actually since the left hand side of (10) is symmetric in $\ell_1, \ell_2, \ell_3, \ell_4$, the right hand side does not depend of the choice of the diagonal.

Note that parity transformation (8) implies that $\ell_1 + \ell_2 + \ell_3 + \ell_4$ must be even and we turn to the field u_ℓ, see 7,

$$\begin{aligned}
&\mathrm{Cum}_4\left(u_{\ell_1}(L_1), u_{\ell_2}(L_2), u_{\ell_3}(L_3), u_{\ell_4}(L_4)\right) \\
&\qquad\qquad = \sum_{m_{1:4}} \prod_{j=1}^{4} Y_{\ell_j}^{m_j}(L_j)\, \mathrm{Cum}_4\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, Z_{\ell_3}^{m_3}, Z_{\ell_4}^{m_4}\right) \\
&= \sum_{\ell^1, m^1} T_4\left(\ell_1, \ell_2, \ell_3, \ell_4 \middle| \ell^1\right) \sqrt{2\ell^1 + 1} \sum_{m_{1:4}} \prod_{j=1}^{4} Y_{\ell_j}^{m_j}(L_j) \begin{pmatrix} \ell_1 & \ell_2 & \ell^1 \\ m_1 & m_2 & -m^1 \end{pmatrix} \begin{pmatrix} \ell^1 & \ell_3 & \ell_4 \\ m^1 & m_3 & m_4 \end{pmatrix}.
\end{aligned}$$

The cumulant in the left hand side is symmetric in $\ell_1, \ell_2, \ell_3, \ell_4$, and invariant under rotation. The function

$$\begin{aligned}
\widetilde{I}_{\ell_1,\ell_2,\ell_3,\ell_4,\ell^1}(L_1, L_2, L_3, L_4) = \sqrt{2\ell^1 + 1} \sum_{m_1,\dots m_4, m^1} \prod_{j=1}^{4} Y_{\ell_j}^{m_j}(L_j) \\
\times \begin{pmatrix} \ell_1 & \ell_2 & \ell^1 \\ m_1 & m_2 & -m^1 \end{pmatrix} \begin{pmatrix} \ell^1 & \ell_3 & \ell_4 \\ m^1 & m_3 & m_4 \end{pmatrix},
\end{aligned}$$

is invariant under the rotation see Lemma 25 Appendix. Therefore we apply the rotation $g_{L_4 L_3}$, which takes the location L_4 to the North pole and at the same time takes L_3 into the plane xOz,

$$\begin{aligned}
&\mathrm{Cum}_4\left(u_{\ell_1}(L_1), u_{\ell_2}(L_2), u_{\ell_3}(L_3), u_{\ell_4}(L_4)\right) \\
&\qquad = \mathrm{Cum}_4\left(u_{\ell_1}(g_{L_4L_3}L_1), u_{\ell_2}(g_{L_4L_3}L_2), u_{\ell_3}(g_{L_4L_3}L_3), u_{\ell_4}(N)\right) \\
&\qquad\quad = \sqrt{\frac{2\ell_4 + 1}{4\pi}} \sum_{\ell^1} T_4\left(\ell_1, \ell_2, \ell_3, \ell_4 \middle| \ell^1\right) \sqrt{2\ell^1 + 1} \\
&\qquad\quad \times \sum_{m_1, m_2, m_3, m^1} \prod_{j=1}^{3} Y_{\ell_j}^{m_j}(g_{L_4L_3}L_j) \begin{pmatrix} \ell_1 & \ell_2 & \ell^1 \\ m_1 & m_2 & -m^1 \end{pmatrix} \begin{pmatrix} \ell^1 & \ell_3 & \ell_4 \\ m^1 & m_3 & 0 \end{pmatrix},
\end{aligned}$$

we have

$$\begin{aligned}
&\mathrm{Cum}_4\left(u_{\ell_1}(L_1), u_{\ell_2}(L_2), u_{\ell_3}(L_3), u_{\ell_4}(L_4)\right) \\
&\qquad\qquad = \sum_{\ell^1} T_4\left(\ell_1, \ell_2, \ell_3, \ell_4 \middle| \ell^1\right) \mathcal{I}_{\ell_1,\ell_2,\ell_3,\ell_4,\ell^1}(\vartheta_1, \vartheta_2, \vartheta_3, \varphi_1, \varphi_2),
\end{aligned}$$

where the function $\mathcal{I}_{\ell_1,\ell_2,\ell_3,\ell_4,\ell^1}$ is given by

$$\mathcal{I}_{\ell_1,\ell_2,\ell_3,\ell_4,\ell^1}\left(\vartheta_1,\vartheta_2,\vartheta_3,\varphi_1,\varphi_2\right)=\sqrt{\frac{2\ell_4+1}{4\pi}\left(2\ell^1+1\right)}\sum_{m_{1:3},m^1}\prod_{j=1}^{3}Y_{\ell_j}^{m_j}\left(g_{L_4L_3}L_j\right)$$
$$\times\begin{pmatrix}\ell_1 & \ell_2 & \ell^1\\ m_1 & m_2 & -m^1\end{pmatrix}\begin{pmatrix}\ell^1 & \ell_3 & \ell_4\\ m^1 & m_3 & 0\end{pmatrix}.$$

Actually $\mathcal{I}_{\ell_1,\ell_2,\ell_3,\ell_4,\ell^1}$ is the rotated version of $\widetilde{I}_{\ell_1,\ell_2,\ell_3,\ell_4,\ell^1}$, it corresponds to the tripolar spherical harmonics defined as irreducible tensor products of the spherical harmonics with different arguments [VMK88], p.160. We define the spherical coordinates $(\vartheta_1,\vartheta_2,\vartheta_3,\varphi_1,\varphi_2)$ as follows $g_{L_4L_3}L_3=(\vartheta_3,0)$, $g_{L_4L_3}L_j=(\vartheta_j,\varphi_j)$, $j=1,2$. Note the orthonormality of the system according to the measure $\prod_{k=1}^{3}\Omega\left(dL_k\right)=\prod_{k=1}^{3}\sin\vartheta_k d\vartheta_k d\varphi_k$, $\vartheta_k\in[0,\pi]$, $\varphi_k\in[0,2\pi]$, i.e.

$$\iiint_{\mathbb{S}_2}\mathcal{I}^*_{\ell_1,\ell_2,\ell_3,\ell_4,\ell^1}\mathcal{I}_{j_1,j_2,j_3,j_4,\ell^2}\prod_{k=1}^{3}\Omega\left(dL_k\right)=\delta_{\ell^1,\ell^2}\prod_{k=1:4}\delta_{\ell_k,j_k},$$

since

$$\left(2\ell_4+1\right)\left(2\ell^1+1\right)\sum_{m^1,m_1,m_2,m_3}\begin{pmatrix}\ell_1 & \ell_2 & \ell^1\\ m_1 & m_2 & -m^1\end{pmatrix}^2\begin{pmatrix}\ell^1 & \ell_3 & \ell_4\\ m^1 & m_3 & 0\end{pmatrix}^2=1,$$

here we sum it up by m_1,m_2 first then by m^1,m_3.

The expression for the $\mathrm{Cum}_4\left(X\left(L_1\right),X\left(L_2\right),X\left(L_3\right),X\left(L_4\right)\right)$ is straightforward

$$\mathrm{Cum}_4\left(X\left(L_1\right),X\left(L_2\right),X\left(L_3\right),X\left(L_4\right)\right)=\sum_{\ell_1,\ell_2,\ell_3,\ell_4=0}^{\infty}\mathrm{Cum}_4\left(u_{\ell_1}\left(L_1\right),\ldots,u_{\ell_4}\left(L_4\right)\right)$$
$$=\sum_{\ell_1,\ldots,\ell_4=0}^{\infty}\sum_{\ell^1}T_4\left(\ell_1,\ell_2,\ell_3,\ell_4\middle|\,\ell^1\right)\mathcal{I}_{\ell_1,\ell_2,\ell_3,\ell_4,\ell^1}\left(L_1,\ldots,L_4\right).$$

Definition 18 *The trispectrum of the isotropic field $X\left(L\right)$ is*

$$S_4\left(\ell_1,\ell_2,\ell_3,\ell_4\middle|\,\ell^1\right)=T_4\left(\ell_1,\ell_2,\ell_3,\ell_4\middle|\,\ell^1\right).$$

In particular, we have formulae for the marginal fourth order cumulants

$$\mathrm{Cum}_4\left(X\left(L_1\right),X\left(L_2\right),X\left(L\right),X\left(L\right)\right)=\sum_{\ell_1,\ell_2,\ell_3,\ell_4=0}^{\infty}\sqrt{\frac{2\ell_3+1}{4\pi}}\sum_{\ell^1}\begin{pmatrix}\ell_3 & \ell_4 & \ell^1\\ 0 & 0 & 0\end{pmatrix}$$
$$\times T_4\left(\ell_1,\ell_2,\ell_3,\ell_4\middle|\,\ell^1\right)\mathcal{I}_{\ell_1,\ell_2,\ell^1}\left(\vartheta_1,\varphi_1,\vartheta_2\right),$$

and

$$
\begin{aligned}
&\operatorname{Cum}_4\left(X\left(L_1\right), X\left(L\right), X\left(L\right), X\left(L\right)\right)\\
&\qquad=\sum_{\ell_1,\ell_2,\ell_3,\ell_4=0}^{\infty}\sqrt{\prod_{j=1}^{4}\frac{2\ell_j+1}{4\pi}}\sum_{\ell^1}T_4\left(\ell_1,\ldots,\ell_4|\,\ell^1\right)\\
&\qquad\qquad\times\sqrt{2\ell^1+1}\begin{pmatrix}\ell_1 & \ell_2 & \ell^1\\ 0 & 0 & 0\end{pmatrix}\begin{pmatrix}\ell^1 & \ell_3 & \ell_4\\ 0 & 0 & 0\end{pmatrix}P_{\ell_1}\left(L\cdot L_1\right)
\end{aligned}
$$

also

$$
\begin{aligned}
&\operatorname{Cum}_4\left(X\left(L\right), X\left(L\right), X\left(L\right), X\left(L\right)\right)\\
&=\sum_{\ell_1,\ell_2,\ell_3,\ell_4=0}^{\infty}\sqrt{\prod_{j=1}^{4}\frac{2\ell_i+1}{4\pi}}\sum_{\ell^1}T_4\left(\ell_1,\ell_2,\ell_3,\ell_4|\,\ell^1\right)\sqrt{2\ell^1+1}\begin{pmatrix}\ell_1 & \ell_2 & \ell^1\\ 0 & 0 & 0\end{pmatrix}\begin{pmatrix}\ell^1 & \ell_3 & \ell_4\\ 0 & 0 & 0\end{pmatrix}.
\end{aligned}
$$

5.2 Linear field

If the triangular array Z_ℓ^m contains independent variables, i.e. the angular momentum field u_ℓ is linear. Then for a fixed degree ℓ

$$
\begin{aligned}
\operatorname{Cum}_4\left(Z_\ell^{m_1}, Z_\ell^{m_2}, Z_\ell^{m_3}, Z_\ell^{m_4}\right) &= \delta_{m_i=m}\sum_{\ell^1,m^1}\begin{pmatrix}\ell_1 & \ell_2 & \ell^1\\ m_1 & m_2 & m^1\end{pmatrix}\begin{pmatrix}\ell^1 & \ell_3 & \ell_4\\ -m^1 & m_3 & m_4\end{pmatrix}\\
&\qquad\times\left(-1\right)^{m^1}\sqrt{2\ell^1+1}T_4\left(\ell,\ell,\ell,\ell|\,\ell^1\right),
\end{aligned}
$$

from the selection rules follows $m^1=-\left(m_1+m_2\right)=-2m$, $m^1=m_3+m_4=2m$, therefore $m_i=0$. Now for similar reason

$$
T_4\left(\ell,\ell,\ell,\ell|\,\ell^1\right)=\sqrt{2\ell^1+1}\operatorname{Cum}_4\left(Z_\ell^0,Z_\ell^0,Z_\ell^0,Z_\ell^0\right)\begin{pmatrix}\ell & \ell & \ell^1\\ 0 & 0 & 0\end{pmatrix}\begin{pmatrix}\ell^1 & \ell & \ell\\ 0 & 0 & 0\end{pmatrix}.
$$

The only nonzero cumulants are

$$
\operatorname{Cum}_4\left(Z_\ell^0,Z_\ell^0,Z_\ell^0,Z_\ell^0\right)=\sum_{\ell^1}\sqrt{2\ell^1+1}\begin{pmatrix}\ell & \ell & \ell^1\\ 0 & 0 & 0\end{pmatrix}^2\begin{pmatrix}\ell^1 & \ell & \ell\\ 0 & 0 & 0\end{pmatrix}^2\operatorname{Cum}_4\left(Z_\ell^0,Z_\ell^0,Z_\ell^0,Z_\ell^0\right),
$$

hence for an isotropic linear field u_ℓ $\operatorname{Cum}_4\left(Z_\ell^{m_1}, Z_\ell^{m_2}, Z_\ell^{m_3}, Z_\ell^{m_4}\right)=0$, for and m_i. If additionally the series of u_ℓ is independent then $\operatorname{Cum}_4\left(X\left(L_1\right), X\left(L_2\right), X\left(L_3\right), X\left(L_4\right)\right)=0$ follows.

6 Higher order spectra for isotropic fields

The generalization of the bispectrum and trispectrum is possible for arbitrary higher order [MP10], [MP11]. If the bispectrum and trispectrum are zero then the field can not be Gaussian, in case all polyspectra are zero then the isotropic field is necessary Gaussian. First we show that the characterization of the isotropy in p^{th} order can be given similarly to the previous cases.

Lemma 19 *The field*

$$X(L) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} Z_\ell^m Y_\ell^m(L),$$

is isotropic in p^{th} *order* ($p > 3$) *iff the cumulant* $\mathrm{Cum}_p\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, \ldots, Z_{\ell_p}^{m_p}\right)$ *of the triangular array* Z_ℓ^m *has the form*

$$\mathrm{Cum}_p\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, \ldots, Z_{\ell_p}^{m_p}\right) = \sum_{\substack{\ell^1,\ldots,\ell^{p-3} \\ m^1,\ldots,m^{p-3}}} (-1)^{\Sigma_{a=1}^{p-3} m^a} \prod_{a=0}^{p-3} \begin{pmatrix} \ell^a & \ell_{a+2} & \ell^{a+1} \\ -m^a & m_{a+2} & m^{a+1} \end{pmatrix} \tag{12}$$

$$\times \prod_{a=1}^{p-3} \sqrt{2\ell^a+1} \widetilde{S}_p\left(\ell_1, \ldots, \ell_p \middle| \ell^1, \ldots, \ell^{p-3}\right),$$

where $\ell^0 = \ell_1$, $\ell^{p-2} = \ell_p$, $k^0 = -k_1$, $k^{p-2} = k_p$, $m^0 = -m_1$, $m^{p-2} = m_p$. *The function* $\widetilde{S}_p\left(\ell_1, \ldots, \ell_p \middle| \ell^1, \ldots, \ell^{p-3}\right)$ *has the form*

$$\widetilde{S}_p\left(\ell_1, \ldots, \ell_p \middle| \ell^1, \ldots, \ell^{p-3}\right) = \sum_{\substack{k_1,\ldots,k_p \\ k^1,\ldots,k^{p-3}}} (-1)^{\Sigma_{a=1}^{p-3} k^a} \prod_{a=0}^{p-3} \begin{pmatrix} \ell^a & \ell_{a+2} & \ell^{a+1} \\ -k^a & k_{a+2} & k^{a+1} \end{pmatrix} \tag{13}$$

$$\times \prod_{a=1}^{p-3} \sqrt{2\ell^a+1}\, \mathrm{Cum}_p\left(Z_{\ell_1}^{k_1}, Z_{\ell_2}^{k_2}, \ldots, Z_{\ell_p}^{k_p}\right).$$

See the Appendix E for the proof. Consider a convex polygon with vertices $A_{1:p-1}$, and edges $\ell_{1:p}$. The diagonals denoted by ℓ^j, $j = 1, 2, \ldots, p-3$, starting from the vertex A_1 divide this polygon into $p-2$ triangles. The first triangle has sides (angular mometum) ℓ_1, ℓ_2 and ℓ^1, the next one has sides ℓ^1, ℓ_3 and ℓ^2, the general one is ℓ^a, ℓ_{a+2} and ℓ^{a+1}, finally the last one ℓ^{p-3}, ℓ_{p-1} and ℓ_p. For each a the sides of the triangle $\left(\ell^a, \ell_{a+2}, \ell^{a+1}\right)$ should fulfil the triangle inequality $\left|\ell^a - \ell^{a+1}\right| \leq \ell_{a+2} \leq \ell^a + \ell^{a+1}$. The coefficients in (12) will differ from zero if orders $-m^a, m_{a+2}, m^{a+1}$, fulfil the assumption $-m^a + m_{a+2} + m^{a+1} = 0$, for all a. This implies $m_1 + m_2 = -m^1$, $m_3 + m^2 = m^1$, $\ldots$, $m_{p-1} + m_p = m^{p-3}$. Let us plug in consecutively m^a and we shall arrive to the result $m_1 + m_2 + \cdots + m_p = 0$. Hence $\mathrm{Cum}_p\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, \ldots, Z_{\ell_p}^{m_p}\right) = 0$, unless $m_1 + m_2 + \cdots + m_p = 0$.

Remark 20 *The cumulant* $\mathrm{Cum}_p\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, \ldots, Z_{\ell_p}^{m_p}\right)$ *is invariant under the order of the quantum numbers* $\ell_1, \ell_2, \ldots, \ell_p$*, hence the right hand side of (12) is invariant as well. The result is that the function* $\widetilde{S}_p\left(\ell_1, \ldots, \ell_p \middle| \ell^1, \ldots, \ell^{p-3}\right)$ *given on values* $\ell_1 \leq \ell_2 \leq \cdots \leq \ell_p$ *will determine all cumulants* $\mathrm{Cum}_p\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, \ldots, Z_{\ell_p}^{m_p}\right)$ *by (12).*

Repeat the representation of the field

$$X\left(L\right) = \sum_{\ell=0}^{\infty} u_\ell\left(L\right),$$

where

$$u_\ell\left(L\right) = \sum_{m=-\ell}^{\ell} Y_\ell^m\left(L\right) Z_\ell^m.$$

We are interested in the invariance of the distribution of $u_\ell\left(L\right)$ under rotations. If the location is fixed then $u_\ell\left(L\right)$ is connected to the distribution of $\mathcal{Z}_\ell^k$, since the Condon and Shortley phase convention (33) provides

$$u_\ell\left(L\right) = \sqrt{\frac{2\ell+1}{4\pi}} \sum_{m=-\ell}^{\ell} D_{m,0}^{(\ell)*} Z_\ell^m.$$

Our main interest is the comparison of the finite dimensional distributions of the process $u_\ell\left(L\right)$ to the rotated one in case the rotation carries one location to the North pole N, since u_ℓ simplifies to

$$u_\ell\left(N\right) = \sqrt{\frac{2\ell+1}{4\pi}} Z_\ell^0.$$

We have

$$\begin{aligned}
\mathrm{Cum}_p\left(u_{\ell_1}\left(L_1\right), u_{\ell_2}\left(L_2\right), \ldots, u_{\ell_p}\left(L_p\right)\right) &= \sum_{m_1,\ldots,m_p} \prod_{j=1}^{p} Y_{\ell_j}^{m_j}\left(L_j\right) \mathrm{Cum}_p\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, \ldots, Z_{\ell_p}^{m_p}\right) \\
= \sum_{m_1,\ldots,m_p} \prod_{j=1}^{p} Y_{\ell_j}^{m_j}\left(L_j\right) &\sum_{\substack{\ell^1,\ldots,\ell^{p-3} \\ m^1,\ldots,m^{p-3}}} (-1)^{\Sigma_{a=1}^{p-3} m^a} \prod_{a=0}^{p-3} \begin{pmatrix} \ell^a & \ell_{a+2} & \ell^{a+1} \\ -m^a & m_{a+2} & m^{a+1} \end{pmatrix} \\
&\times \prod_{a=1}^{p-3} \sqrt{2\ell^a+1} \widetilde{S}_p\left(\ell_1, \ldots, \ell_p \middle| \ell^1, \ldots, \ell^{p-3}\right) \\
&= \sum_{\ell^1,\ldots,\ell^{p-3}} \widetilde{I}_{\ell_{1:p},\ell^{1:p-3}}\left(L_1, \ldots, L_p\right) \widetilde{S}_p\left(\ell_1, \ldots, \ell_p \middle| \ell^1, \ldots, \ell^{p-3}\right).
\end{aligned}$$

The following $p-$product *of spherical harmonics* Y_ℓ^m

$$\begin{aligned}\widetilde{I}_{\ell_1,\ldots,\ell_p,\ell^1,\ldots,\ell^{p-3}}\left(L_1,\ldots,L_p\right) &= \sum_{\substack{m_1,\ldots,m_p\\ m^1,\ldots,m^{p-3}}} \prod_{j=1}^{p} Y_{\ell_j}^{m_j}\left(L_j\right)\\ &\quad\times\left(-1\right)^{\Sigma k^{1:p-3}} \prod_{a=0}^{p-3}\begin{pmatrix}\ell^a & \ell_{a+2} & \ell^{a+1}\\ -m^a & m_{a+2} & m^{a+1}\end{pmatrix}\prod_{a=1}^{p-3}\sqrt{2\ell^a+1},\end{aligned}$$

is rotation invariant, see (30), hence we can apply the rotation $g_{L_pL_{p-1}}$ such that $g_{L_pL_{p-1}}L_p = N$, and $g_{L_pL_{p-1}}L_{p-1}$ belongs into the plane xOz, we have

$$\begin{aligned}I_{\ell_1,\ldots,\ell_p,\ell^1,\ldots,\ell^{p-3}}\left(L_1,\ldots,L_p\right) &= \sqrt{\frac{2\ell_p+1}{4\pi}} \sum_{\substack{m_1,\ldots,m_p\\ m^1,\ldots,m^{p-3}}} \prod_{j=1}^{p-1} Y_{\ell_j}^{m_j}\left(g_{L_pL_{p-1}}L_j\right)\\ \times\left(-1\right)^{\Sigma k^{1:p-3}} \prod_{a=0}^{p-3}\begin{pmatrix}\ell^a & \ell_{a+2} & \ell^{a+1}\\ -m^a & m_{a+2} & m^{a+1}\end{pmatrix}&\prod_{a=1}^{p-3}\sqrt{2\ell^a+1}\begin{pmatrix}\ell^{p-3} & \ell_{p-1} & \ell_p\\ -m^{p-3} & m_{p-1} & 0\end{pmatrix}\sqrt{2\ell_p+1}.\end{aligned}$$

According to the usual measure $\prod_{k=1}^{p-1}\Omega\left(dL_k\right) = \prod_{k=1}^{p-1}\sin\vartheta_k d\vartheta_k d\varphi_k$, $\vartheta_k \in [0,\pi]$, $\varphi_k \in [0,2\pi]$, the system of functions $I_{\ell_1,\ldots,\ell_p,\ell^1,\ldots,\ell^{p-3}}$ forms an orthogonal system.

Consider now

$$\begin{aligned}\widetilde{S}_p\left(\ell_1,\ldots,\ell_p|\,\ell^1,\ldots,\ell^{p-3}\right) &= \sum_{\substack{m_1,\ldots,m_p\\ m^1,\ldots,m^{p-3}}} \left(-1\right)^{\Sigma_{a=1}^{p-3}m^a}\prod_{a=0}^{p-3}\begin{pmatrix}\ell^a & \ell_{a+2} & \ell^{a+1}\\ m^a & m_{a+2} & m^{a+1}\end{pmatrix}\\ &\quad\times\prod_{a=1}^{p-3}\sqrt{2\ell^a+1}\,\mathrm{Cum}_p\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2},\ldots,Z_{\ell_p}^{m_p}\right)\end{aligned}$$

where remember $\ell^0 = \ell_1$, and $\ell^{p-2} = \ell_p$. The left size is symmetric in $\ell_1,\ldots,\ell_p$ hence the right size must be symmetric as well. One might fix an order for the entries of $\ell_1,\ldots,\ell_p$ to get a unique representation for the cumulant. We consider a monotone ordering $\ell_1 \le \ell_2 \le \ldots \le \ell_p$ and refer to it as canonical representation.

Note that parity transformation (8) implies that $\ell_1+\ell_2+\ell_3+\ldots+\ell_p$ must be even.

$$\begin{aligned}\mathrm{Cum}_p\left(X\left(L_1\right),X\left(L_2\right),X\left(L_3\right),\ldots,X\left(L_p\right)\right) &= \sum_{\ell_{1:p}=0}^{\infty}\mathrm{Cum}_p\left(u_{\ell_1}\left(L_1\right),u_{\ell_2}\left(L_2\right),\ldots,u_{\ell_p}\left(L_p\right)\right)\\ &= \sum_{\ell_{1:p}=0}^{\infty}\widetilde{S}_p\left(\ell_1,\ldots,\ell_p|\,\ell^1,\ldots,\ell^{p-3}\right)I_{\ell_1,\ldots,\ell_p,\ell^1,\ldots,\ell^{p-3}}\left(L_1,\ldots,L_p\right).\end{aligned}$$

Definition 21 *The* p^{th} *order polyspectrum of the isotropic field* $X\left(L\right)$ *is* $S_p\left(\ell_1,\ldots,\ell_p|\,\ell^1,\ldots,\ell^{p-3}\right) = \widetilde{S}_p\left(\ell_1,\ldots,\ell_p|\,\ell^1,\ldots,\ell^{p-3}\right)$.

6.1 Linear field

Let us consider the particular case when Z_ℓ^m are independent if ℓ is fixed. We have

$$\operatorname{Cum}_p\left(Z_\ell^{m_1}, Z_\ell^{m_2}, \ldots, Z_\ell^{m_p}\right) = \delta_{m_i=m} \sum_{\substack{\ell^1,\ldots,\ell^{p-3}\\ m^1,\ldots,m^{p-3}}} (-1)^{\Sigma_{a=1}^{p-3} m^a} \prod_{a=0}^{p-3} \begin{pmatrix} \ell^a & \ell & \ell^{a+1} \\ -m^a & m_{a+2} & m^{a+1} \end{pmatrix}$$

$$\times S_p\left(\ell \middle| \ell^1, \ldots, \ell^{p-3}\right) \prod_{a=1}^{p-3} \sqrt{2\ell^a + 1}.$$

We have seen that for $p = 3, 4$, all $m_i = 0$ follows. Let us see the case $p = 5$,

$$\begin{pmatrix} \ell & \ell^1 \\ m_{1:2} & m^1 \end{pmatrix} \begin{pmatrix} \ell^1 & \ell & \ell^2 \\ -m^1 & m_3 & m^2 \end{pmatrix} \begin{pmatrix} \ell^2 & \ell & \ell \\ -m^2 & m_4 & m_5 \end{pmatrix}$$

then from the selection rules follows that $m^1 = -(m_1 + m_2) = -2m$, $m^2 = m^1 - m_3 = -3m$, $m^2 = m_4 + m_5 = 2m$, hence $m = 0$. In general it is easy to see that $m^k = -(k+1)m$, for $k = 1, 2, \ldots, p-3$, and at the same time $m^{p-3} = 2m$, hence $m = 0$, and $m^j = 0$ as well.

Consider the polyspectrum

$$\widetilde{S}_p\left(\ell \middle| \ell^{1:p-3}\right) = \sum_{\substack{k_1,\ldots,k_p\\ k^1,\ldots,k^{p-3}}} (-1)^{\Sigma k^{1:p-2}} \prod_{a=0}^{p-3} \begin{pmatrix} \ell^a & \ell & \ell^{a+1} \\ -k^a & k_{a+2} & k^{a+1} \end{pmatrix} \prod_{a=1}^{p-3} \sqrt{2\ell^a + 1}\, \operatorname{Cum}_p\left(Z_\ell^{k_{1:p}}\right),$$

from the independence follows that $\ell_j = \ell$ and $k_j = k$, for $j = 1, 2, \ldots, p$. Hence similar argument to the previous one leads to the result: $k_j = 0$, and $k^j = 0$ as well. We have

$$\widetilde{S}_p\left(\ell \middle| \ell^{1:p-3}\right) = \prod_{a=0}^{p-3} \begin{pmatrix} \ell^a & \ell & \ell^{a+1} \\ 0 & 0 & 0 \end{pmatrix} \prod_{a=1}^{p-3} \sqrt{2\ell^a + 1}\, \operatorname{Cum}_p\left(Z_\ell^0, Z_\ell^0, \ldots, Z_\ell^0\right).$$

Now, by the Lemma 19 we get

$$\operatorname{Cum}_p\left(Z_\ell^0, Z_\ell^0, \ldots, Z_\ell^0\right) = \sum_{\ell^1,\ldots,\ell^{p-3}} \prod_{a=0}^{p-3} \begin{pmatrix} \ell^a & \ell & \ell^{a+1} \\ 0 & 0 & 0 \end{pmatrix}^2 \prod_{a=1}^{p-3} \sqrt{2\ell^a + 1}\, \operatorname{Cum}_p\left(Z_\ell^0, Z_\ell^0, \ldots, Z_\ell^0\right).$$

Hence $\operatorname{Cum}_p\left(Z_{\ell_1}^{k_1}, Z_{\ell_2}^{k_2}, \ldots, Z_{\ell_p}^{k_p}\right) = 0$, for all k_j. Now we have a general conclusion because the only case when all cumulants vanish except the second order one is the Gaussian. Once the rows of $\{Z_\ell^m\}$ are Gaussian then all the entries of $\{Z_\ell^m\}$ are independent. Indeed the isotropy implies that all the entries of $\{Z_\ell^m\}$ are uncorraleted now they are Gaussian hence they are independent.

Lemma 22 *If the isotropic field* $X(L)$ *is linear, i.e. inside the rows of the generating array* Z_ℓ^m *all random variables are independent, then the whole array contains independent entries and* $X(L)$ *is Gaussian.*

7 Construction of isotropic field

The stochastic isotropic field

$$X\left(L\right)=\sum_{\ell=0}^{\infty}\sum_{m=-\ell}^{\ell}Z_{\ell}^{m}Y_{\ell}^{m}\left(L\right),$$

on sphere has very spacial angular spectra, in particular the form of the cumulants of the triangular array $\{Z_{\ell}^{m}\}$ is restricted, see (12). If one starts with a non-Gaussian continuos in mean square $X\left(L\right)$ then the triangular array $\{Z_{\ell}^{m}\}$ naturally given by the inversion formula

$$Z_{\ell}^{m}=\int_{\mathbb{S}_2}X\left(L\right)Y_{\ell}^{m*}\left(L\right)dL.$$

Now, we address the reverse question of construction of a triangular array $\{Z_{\ell}^{m}\}$ with the desired properties of their cumulants. Let us start with triangular array $\left\{\widetilde{Z}_{\ell}^{m}\right\}$, assume it is uncorrelated and all moments exist. Consider the vectors $\widetilde{Z}_{\ell}=\left[\widetilde{Z}_{\ell}^{-\ell},\widetilde{Z}_{\ell}^{-\ell+1},\ldots,\widetilde{Z}_{\ell}^{\ell}\right]^{\top}$, $.\ell=0,1,2,\ldots$, according to the rows of $\left\{\widetilde{Z}_{\ell}^{m}\right\}$. The finite dimensional distribution is characterized by its cumulant function (logarithm of the characteristic function)

$$\Phi_{\widetilde{Z}}\left(\omega_{\ell}\middle|\,\ell=0,1,2,\ldots\right)=\ln\mathsf{E}\exp\left(i\sum_{\ell}\omega_{\ell}^{\top}\widetilde{Z}_{\ell}\right),$$

such that only finite many coordinates of the variables $\left(\omega_{\ell}\middle|\,\ell=0,1,2,\ldots\right)$ are different from zero. Consider the transformed series $\widehat{Z}_{\ell}=D^{(\ell)}\widetilde{Z}_{\ell}$, where $D^{(\ell)}$ is the Wigner matrix of rotations $D^{(\ell)}=\left[D_{k,m}^{(\ell)}\right]_{k,m=-\ell}^{\ell}$, and define a triangular array $\{Z_{\ell}^{m}\}$ through the cumulant function

$$\Phi_{Z}\left(\omega_{\ell}\middle|\,\ell=0,1,2,\ldots\right)=\int_{SO(3)}\ln\mathsf{E}\exp\left(i\sum_{\ell}\omega_{\ell}^{\top}D^{(\ell)}\widetilde{Z}_{\ell}\right)dg$$

where again only finite many coordinates of the variables $\left(\omega_{\ell}\middle|\,\ell=0,1,2,\ldots\right)$ are different from zero and $dg=\sin\vartheta d\vartheta d\varphi d\gamma/8\pi^2$, is the Haar measure with unite mass. This new triangular array $\{Z_{\ell}^{m}\}$ will be called Wigner D-transform of $\left\{\widetilde{Z}_{\ell}^{m}\right\}$. .The third order cumulants of Z_{ℓ}^{m} for instance

$$\begin{aligned}\operatorname{Cum}_3\left(Z_{\ell_1}^{k_1},Z_{\ell_2}^{k_2},Z_{\ell_3}^{k_3}\right)&=\int_{SO(3)}\sum_{m_1,m_2,m_3}D_{k_1,m_1}^{(\ell_1)}D_{k_2,m_2}^{(\ell_2)}D_{k_3,m_3}^{(\ell_3)}dg\operatorname{Cum}_3\left(\widetilde{Z}_{\ell_1}^{m_1},\widetilde{Z}_{\ell_2}^{m_2},\widetilde{Z}_{\ell_3}^{m_3}\right)\\&=\begin{pmatrix}\ell_1&\ell_2&\ell_3\\k_1&k_2&k_3\end{pmatrix}\sum_{m_1,m_2,m_3}\begin{pmatrix}\ell_1&\ell_2&\ell_3\\m_1&m_2&m_3\end{pmatrix}\operatorname{Cum}_3\left(\widetilde{Z}_{\ell_1}^{m_1},\widetilde{Z}_{\ell_2}^{m_2},\widetilde{Z}_{\ell_3}^{m_3}\right)\\&=\begin{pmatrix}\ell_1&\ell_2&\ell_3\\k_1&k_2&k_3\end{pmatrix}B_3\left(\ell_1,\ell_2,\ell_3\right),\end{aligned}$$

which fulfils (9). The function $B_3(\ell_1, \ell_2, \ell_3)$ is given by

$$B_3(\ell_1, \ell_2, \ell_3) = \sum_{m_1, m_2, m_3} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \operatorname{Cum}_3\left(\widetilde{Z}_{\ell_1}^{m_1}, \widetilde{Z}_{\ell_2}^{m_2}, \widetilde{Z}_{\ell_3}^{m_3}\right),$$

and also

$$B_3(\ell_1, \ell_2, \ell_3) = \sum_{m_1, m_2, m_3} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \operatorname{Cum}_3\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, Z_{\ell_3}^{m_3}\right).$$

The conclusion is that a subset of cumulants $\operatorname{Cum}_3\left(\widetilde{Z}_{\ell_1}^{m_1}, \widetilde{Z}_{\ell_2}^{m_2}, \widetilde{Z}_{\ell_3}^{m_3}\right)$, i.e. $m_1 + m_2 + m_3 = 0$, is used in the construction and the superposition with 'probability' amplitudes is applied. In general we also have

$$\begin{aligned} \operatorname{Cum}_p\left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, \dots, Z_{\ell_p}^{m_p}\right) = & \sum_{\substack{\ell^1, \dots, \ell^{p-3} \\ m^1, \dots, m^{p-3}}} (-1)^{\Sigma_{a=1}^{p-3} m^a} \prod_{a=0}^{p-3} \begin{pmatrix} \ell^a & \ell_{a+2} & \ell^{a+1} \\ -m^a & m_{a+2} & m^{a+1} \end{pmatrix} \\ & \times \prod_{a=1}^{p-3} \sqrt{2\ell^a + 1} S_p\left(\ell_{1:p} \middle| \ell^{1:p-3}\right) \end{aligned}$$

where

$$\begin{aligned} S_p\left(\ell_{1:p} \middle| \ell^{1:p-3}\right) = & \sum_{\substack{m_1, \dots, m_p \\ m^1, \dots, m^{p-3}}} (-1)^{\Sigma_{a=1}^{p-3} m^a} \prod_{a=0}^{p-3} \begin{pmatrix} \ell^a & \ell_{a+2} & \ell^{a+1} \\ -m^a & m_{a+2} & m^{a+1} \end{pmatrix} \\ & \times \prod_{a=1}^{p-3} \sqrt{2\ell^a + 1} \operatorname{Cum}_p\left(\widetilde{Z}_{\ell_1}^{k_1}, \widetilde{Z}_{\ell_2}^{k_2}, \dots, \widetilde{Z}_{\ell_p}^{k_p}\right). \end{aligned}$$

A Cumulants

A.1 Basic properties

Cumulants (also called invariants) are very important quantities for the characterization of a random series, see [RST06] for some details. Moments and cumulants are equvivalent, since the moments can be defined as the coefficients in the series expansion of the characteristic function similarly cumulants are the coefficients in the series expansion of the log-characteristic function. The main importance of the cumulants is that it does not contain the Gaussian part of the moments. Let $\underline{X} \in R^n$ denote a random vector.

A.1.1 Symmetry

$$\operatorname{Cum}(\underline{X}) = \operatorname{Cum}(\underline{X}_{\mathfrak{p}(1:n)}), \ \underline{X} \in \mathbb{R}^n,$$

where $\mathfrak{p}(1:n) = (\mathfrak{p}(1), \mathfrak{p}(2), \dots, \mathfrak{p}(n))$, $p \in \mathfrak{P}_\mathfrak{n}$ and $\mathfrak{P}_\mathfrak{n}$ denotes the set of all permutations of the integers $(1:n)$.

A.1.2 Multilinearity

For any constants $c_{1:2} = (c_1, c_2)$

$$\mathrm{Cum}(c_1 Y_1 + c_2 Y_2, \underline{X}) = c_1 \,\mathrm{Cum}(Y_1, \underline{X}) + c_2 \,\mathrm{Cum}(Y_2, \underline{X}).$$

A.1.3 Independence

If $\underline{X} \in R^n$ is independent of $\underline{Y} \in R^m$ where $n, m > 0$ then

$$\mathrm{Cum}(\underline{X}, \underline{Y}) = 0.$$

In particular if $n = m$ then for independent $\underline{X}$ and $\underline{Y}$

$$\mathrm{Cum}(\underline{X} + \underline{Y}) = \mathrm{Cum}(\underline{X}) + \mathrm{Cum}(\underline{Y}).$$

This formula is the additive version of the formula of the moment of the product of independent variables.

A.1.4 Gaussianity

The random vector $\underline{X} \in R^n$ is Gaussian if and only if for all vector $k_{(1:m)}$ with elements from $(1 : n)$

$$\mathrm{Cum}(\underline{X}_{k_{(1:m)}}) = 0, \;\; m > 2.$$

A.1.5 Expression of the cumulant via moments

The first order cumulants is the moment

$$\mathrm{Cum}(X) = \mathsf{E}X,$$

the second and third order cumulants are the second and third order central moments

$$\mathrm{Cum}(X_1, X_2) = \mathrm{Cov}(X_1, X_2) \tag{14}$$

$$\mathrm{Cum}(X_1, X_2, X_3) = \mathsf{E}\prod_{i=1}^{3} (X_i - \mathsf{E}X_i)\,. \tag{15}$$

The general formula is

$$\mathrm{Cum}(\underline{X}) = \sum_{m=1}^{n} (-1)^{m-1}(m-1)! \sum_{U \in \mathcal{P}_n} \prod_{j=1}^{m} \mathsf{E}\; X_{1:n}^{u_{j,1:n}}, \;\; \underline{X} \in \mathbb{R}^n, \tag{16}$$

where the second summation is taken over all possible partitions U, where $[u_{j,1:n}]_{j=1:m}$ is an indicator of a partition $\mathcal{L}$ with m subsets of the set $(1 : n) = (1, 2, \ldots, n)$, $u_{j,k} = 1$ if $k \in \mathcal{L}$, otherwise 0. The double sum in (16) is over all partitions of $\mathcal{L} \in \mathcal{P}_n$, where $\mathcal{P}_n$ is the set of all partitions of the numbers $(1 : n)$.

Example 23 *First note that if $n = 3$, then the m^{th} order partitions $\mathcal{L} \in \mathcal{P}_3$ with the corresponding indicators and permutations of $(1:3)$ are*

1. *for $m = 1$; $\mathcal{L} = \{(1:3)\}$*

$$\mathcal{L} \leftrightarrow u_{1,1:3} = [1,1,1], \quad \mathsf{E}\ X_{1:3}^{u_{1,1:3}} = \mathsf{E}X_1X_2X_3,$$

2. *for $m = 2$; $\mathcal{L}_1 = \{(1),\ (2,3)\}$, $\mathcal{L}_2 = \{(1,3),\ (2)\}$, $\mathcal{L}_3 = \{(1,2),\ (3)\}$,*

$$\mathcal{L}_1 \leftrightarrow \begin{bmatrix} u_{1,1:3} \\ u_{2,1:3} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix}, \quad \prod_{j=1}^{2} \mathsf{E}\ X_{1:3}^{u_{j,1:3}} = \mathsf{E}X_1\mathsf{E}X_2X_3,$$

$$\mathcal{L}_2 \leftrightarrow \begin{bmatrix} u_{1,1:3} \\ u_{2,1:3} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, \quad \prod_{j=1}^{2} \mathsf{E}\ X_{1:3}^{u_{j,1:3}} = \mathsf{E}X_2\mathsf{E}X_1X_3,$$

$$\mathcal{L}_3 \leftrightarrow \begin{bmatrix} u_{1,1:3} \\ u_{2,1:3} \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad \prod_{j=1}^{2} \mathsf{E}\ X_{1:3}^{u_{j,1:3}} = \mathsf{E}X_3\mathsf{E}X_1X_2,$$

3. *for $m = 3$; $\mathcal{L} = \{(1),\ (2),(3)\}$*

$$\mathcal{L} \leftrightarrow \begin{bmatrix} u_{1,1:3} \\ u_{2,1:3} \\ u_{2,1:3} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad \prod_{j=1}^{3} \mathsf{E}\ X_{1:3}^{u_{j,1:3}} = \mathsf{E}X_1\mathsf{E}X_2\mathsf{E}X_3,$$

Now we apply (16):

$$\begin{aligned} \operatorname{Cum}(X_1,X_2,X_3) &= \mathsf{E}X_1X_2X_3 - \mathsf{E}X_1\mathsf{E}X_2X_3 - \mathsf{E}X_2\mathsf{E}X_1X_3 \\ &\quad - \mathsf{E}X_3\mathsf{E}X_1X_2 + 2\mathsf{E}X_1\mathsf{E}X_2\mathsf{E}X_3 \\ &= \mathsf{E}\prod_{i=1}^{3}(X_i - \mathsf{E}X_i), \end{aligned}$$

The first three cumulants, see also (14), equal the central moments but this is not true for higher order cumulants. One might easily check this for the case of cumulants of order four. Indeed, $\underline{X} \in \mathbb{R}^4$

$$\begin{aligned} \operatorname{Cum}(\underline{X}) &= \operatorname{Cum}(\underline{X} - \mathsf{E}\underline{X}) \\ &= \mathsf{E}\prod_{i=1}^{4}(X_i - \mathsf{E}X_i) - \operatorname{Cov}(X_1,X_2)\operatorname{Cov}(X_3,X_4) \\ &\quad - \operatorname{Cov}(X_1,X_3)\operatorname{Cov}(X_2,X_4) - \operatorname{Cov}(X_1,X_4)\operatorname{Cov}(X_2,X_3) \\ &\neq \mathsf{E}\prod_{i=1}^{4}(X_i - \mathsf{E}\ X_i), \end{aligned} \tag{17}$$

unless all covariances are zero.

A.1.6 Expression of the moment via cumulants

We consider the moment $\mathsf{E}\prod_{i=1}^{n} X_i$ as the general case because the moment $\mathsf{E}Y_{1:m}^{k_{1:m}}$ can be put into the form $\mathsf{E}\prod_{i=1}^{n} X_i$, where we define $n = \Sigma k_{(1:m)}$ and

$$\underline{X} = (\underbrace{Y_1, \ldots, Y_1}_{k_1}, \ldots, \underbrace{Y_m, \ldots, Y_m}_{k_m}).$$

Now

$$\mathsf{E}\prod_{i=1}^{n} X_i = \sum_{\mathcal{L}\in\mathcal{P}_n} \prod_{\mathbf{b}_j\in\mathcal{L}} \operatorname{Cum}(X_{\mathbf{b}_j}), \tag{18}$$

where the summation is over all partitions $\mathcal{L} = \{\mathbf{b}_1, \mathbf{b}_2, \ldots, \mathbf{b}_k\}$ of $(1:n)$.

An example of the use of the formula (18) can be seen when $n = 3$.

Example 24 *Each partition of* $(1:3)$ *are listed in the previus example therefore if* $\underline{X} \in \mathbb{R}^3$,

$$\begin{aligned} \mathsf{E}X_1X_2X_3 = {} & \operatorname{Cum}(X_{1:3}) + \operatorname{Cum}(X_1)\operatorname{Cum}(X_2, X_3) \\ & + \operatorname{Cum}(X_2)\operatorname{Cum}(X_1, X_3) + \operatorname{Cum}(X_3)\operatorname{Cum}(X_1, X_2) \\ & + \operatorname{Cum}(X_1)\operatorname{Cum}(X_2)\operatorname{Cum}(X_3). \end{aligned}$$

Now in particular

$$\mathsf{E}\ X^3 = \operatorname{Cum}(X, X, X) + 3\operatorname{Cum}(X)\operatorname{Cum}(X, X) + \operatorname{Cum}(X)^3.$$

B Spherical Harmonics

The following notation and results are used in this mauscript.

1. $\mathbb{R}^3$ is Euclidean space of dimension 3.
2. $\mathbb{S}_2$: **sphere** with radius 1 in $\mathbb{R}^3$, colatitude (angular) coordinate $\vartheta \in [0, \pi]$, longitude coordinate $\varphi \in [0, 2\pi]$, North pole N: $\vartheta = 0$, φ is indeterminate, (latitude coordinate $\pi/2 - \vartheta \in [-\pi/2, \pi/2]$, North pole: $\vartheta = \pi/2$,) The spherical coordinates on $\mathbb{S}_2$: $(\sin\vartheta\cos\varphi, \sin\vartheta\sin\varphi, \cos\vartheta)$
3. $SO(3)$ denotes the 3D (special orthogonal) rotation group
4. **Addition Theorem**. [EMOT81] vol. 2,7.15 (30), pp.116 Let U_1 and U_2 be two vectors with angle ϑ and lengths r_1 and r_2 respectively, and the Euclidean distance is denoted by $\rho = \|U_1 - U_2\| = \sqrt{r_1^2 + r_2^2 - 2r_1r_2\cos\gamma}$, then we have

$$\rho^{-\nu} J_\nu(\rho) = \left(\frac{r_1r_2}{2}\right)^{-\nu} \Gamma(\nu) \sum_{\ell=0}^{\infty} (\ell + \nu)\, C_\ell^\nu(\cos\gamma)\, J_{\ell+\nu}(r_1)\, J_{\ell+\nu}(r_2), \tag{19}$$

where J_ν denotes the Bessel function of the first kind, C_ℓ^ν Gegenbauer polynomial $\left(C_\ell^{1/2} = P_\ell\right)$.

5. **Lapalce-Beltrami operator** $\triangle_B = \triangledown_B^2$,

$$\triangle_B = \frac{1}{\sin\vartheta}\frac{\partial}{\partial\vartheta}\left(\sin\vartheta\frac{\partial}{\partial\vartheta}\right) + \frac{1}{\sin^2\vartheta}\frac{\partial^2}{\partial\varphi^2},$$
$$\triangle_B = \frac{1}{\sin\vartheta}\left[\frac{\partial}{\partial\vartheta}\left(\sin\vartheta\frac{\partial}{\partial\vartheta}\right) + \frac{1}{\sin\vartheta}\frac{\partial^2}{\partial\varphi^2}\right],$$
$$\triangle_B = \frac{\partial^2}{\partial\vartheta^2} + \frac{\cos\vartheta}{\sin\vartheta}\frac{\partial}{\partial\vartheta} + \frac{1}{\sin^2\vartheta}\frac{\partial^2}{\partial\varphi^2},$$

([Apo07], vol. 2 pp. 293).

6. $\triangledown$ is the central difference operator $\left(\triangledown_1^2 + \triangledown_2^2\right) X_{j,k} = X_{j+1,k} + X_{j-1,k} + X_{j,k+1} + X_{j,k-1} - 4X_{j,k}$

7. **Spherical function** f of order ℓ.

$$\begin{aligned}\triangle_B f &= \frac{1}{\sin\vartheta}\frac{\partial}{\partial\vartheta}\left(\sin\vartheta\frac{\partial f}{\partial\vartheta}\right) + \frac{1}{\sin^2\vartheta}\frac{\partial^2 f}{\partial\varphi^2}\\ &= -\ell\left(\ell+1\right) f,\end{aligned}$$

[VMK88]. *Regular* **spherical harmonics** Y_ℓ, homogeneous polynomial of degree ℓ and

$$\nabla^2 Y_\ell = 0.$$

8. *Normalized* **Legendre polynomial** $\widetilde{P}_\ell$ (*Rodrigues' formula*)

$$\widetilde{P}_\ell\left(x\right) = \sqrt{\frac{2\ell+1}{2}}\frac{1}{2^\ell \ell!}\frac{d^\ell\left(x^2-1\right)^\ell}{dx^\ell},\ x\in\left[-1,1\right],$$
$$\int_{-1}^{1}\left(\widetilde{P}_\ell\left(x\right)\right)^2 dx = 1$$

Standardized **Legendre polynomial** $P_0\left(x\right) = 1$,

$$P_\ell\left(x\right) = \frac{1}{2^\ell \ell!}\frac{d^\ell\left(x^2-1\right)^\ell}{dx^\ell},\ x\in\left[-1,1\right],$$

$P_\ell\left(1\right) = 1$([EMOT81], vol. 2, pp.180) it is orthogonal and

$$\int_{-1}^{1}\left[P_\ell\left(x\right)\right]^2 dx = \frac{2}{2\ell+1}$$

$$\begin{aligned}\int_{\mathbb{S}_2}\left[P_\ell\left(\cos\vartheta\right)\right]^2\Omega\left(dL\right) &= \int_0^{2\pi}\int_0^{\pi}\left[P_\ell\left(\cos\vartheta\right)\right]^2\sin\vartheta d\vartheta d\varphi\\ &= 2\pi\int_0^{\pi}\left[P_\ell\left(\cos\vartheta\right)\right]^2\sin\vartheta d\vartheta\\ &= 2\pi\int_{-1}^{1}\left[P_\ell\left(x\right)\right]^2 dx\\ &= \frac{4\pi}{2\ell+1}\end{aligned} \tag{20}$$

recurrence

$$(\ell+1)\,P_{\ell+1}(x) = (2\ell+1)\,xP_{\ell}(x) - \ell P_{\ell-1}(x)$$

generating functions ([EMOT81], vol. 2, pp.183)

$$\sum_{\ell=0}^{\infty} P_{\ell}(x)\,z^{\ell} = \left(1-2xz+z^{2}\right)^{-1/2},\ x\in(-1,1),\ |z|<1 \tag{21}$$

$$\sum_{\ell=0}^{\infty} \frac{1}{\ell!}P_{\ell}(\cos\gamma)\,z^{\ell} = e^{z\cos\gamma}J_0(z\sin\vartheta) \tag{22}$$

$$\sum_{\ell=0}^{\infty} \frac{(-1)^{\ell}}{\ell+1/2}P_{\ell}(\cos\gamma)\,z^{2\ell+1} = F(\sin\vartheta/2,\varphi), \quad z=\tan\frac{\varphi}{2}, \quad 0<\varphi<\frac{\pi}{2}, 0<\vartheta<\pi.$$

9. P_{ℓ}^{m} is the *associated normalized* **Legendre function** *of the first kind* (Gegenbauer polynomial at particular indices) of **degree** ℓ and **order** m

$$P_{\ell}^{m}(x) = (-1)^{m}\left(1-x^{2}\right)^{m/2}\frac{d^{m}P_{\ell}(x)}{dx^{m}},$$
$$P_{\ell}^{m}(z) = \left(z^{2}-1\right)^{m/2}\frac{d^{m}P_{\ell}(z)}{dz^{m}},$$

recurrence formula is

$$(\ell-m+1)\,P_{\ell+1}^{m}(x) = (2\ell+1)\,xP_{\ell}^{m}(x) - (\ell+m)\,P_{\ell-1}^{m}(x),$$

in particular $P_{\ell}^{m}(1)=\delta_{m,0}$, $.P_{\ell}^{0}(x)=P_{\ell}(x)$,

$$P_{\ell}^{-m}(x) = (-1)^{m}\frac{\Gamma(\ell-m+1)}{\Gamma(\ell+m+1)}P_{\ell}^{m}(x).$$

10. **Funk-Hecke formula** ([Ml66]) Suppose G is continuous on $[-1,1]$, then for any spherical harmonic $Y_{\ell}(L)$

$$\int_{\mathbb{S}_2} G(L_1\cdot L)\,Y_{\ell}(L)\,\Omega(dL) = cY_{\ell}(L_1),$$
$$c = 2\pi\int_{-1}^{1} G(x)\,P_{\ell}(x)\,dx,$$

where $\Omega(dL)=\sin\vartheta d\vartheta d\varphi$ is Lebesque element of surface area on $\mathbb{S}_2$, $L_1\cdot L_2=\cos\vartheta$. In particular

$$\int_{\mathbb{S}_2} G(L_1\cdot L)\,Y_{\ell}^{m}(L)\,\Omega(dL) = g_{\ell}Y_{\ell}^{m}(L_1), \tag{23}$$
$$g_{\ell} = 2\pi\int_{-1}^{1} G(x)\,P_{\ell}(x)\,dx$$

see [Yad83] pp.72, for a general dimension d instead of P_ℓ one has the Gegenbauer polynomial $C_\ell^{\frac{d-2}{2}}$ $\left(C_\ell^{1/2} = P_\ell\right)$.
Also

$$\int_{\mathbb{S}_2} G\left(L_1 \cdot L\right) P_\ell\left(L_2 \cdot L\right) \Omega\left(dL\right) = f_\ell P_\ell\left(L_1 \cdot L_2\right)$$

$$f_\ell = 2\pi \int_{-1}^{1} G\left(x\right) P_\ell\left(x\right) dx$$

here $f_\ell = \left(LT\right)_\ell G$ is the Legendre transform of G. Hence

$$G\left(\cos\vartheta\right) = \sum_{\ell=0}^{\infty} \frac{2\ell+1}{2} \int_{-1}^{1} G\left(x\right) P_\ell\left(x\right) dx P_\ell\left(\cos\vartheta\right)$$

In particular *Funk-Hecke formula when* N denotes the north pole

$$\int_{\mathbb{S}_2} \left[P_\ell\left(N \cdot L\right)\right]^2 \Omega\left(dL\right) = cP_\ell\left(N \cdot N\right)$$

$$c = 2\pi \int_{-1}^{1} \left[P_\ell\left(x\right)\right]^2 dx, \quad P_\ell\left(1\right) = 1$$

$$\int_{\mathbb{S}_2} \left[P_\ell\left(N \cdot L\right)\right]^2 \Omega\left(dL\right) = \frac{4\pi}{2\ell+1}$$

11. **Orthonormal spherical harmonics with complex values** $Y_\ell^m\left(\vartheta,\varphi\right)$, $\ell = 0, 1, 2, \ldots$, $m = -\ell, -\ell+1, \ldots -1, 0, 1, \ldots, \ell-1, \ell$ of **degree** ℓ and **order** m (rank ℓ and projection m)

$$Y_\ell^m\left(\vartheta,\varphi\right) = \left(-1\right)^m \sqrt{\frac{2\ell+1}{4\pi}\frac{\left(\ell-m\right)!}{\left(\ell+m\right)!}} P_\ell^m\left(\cos\vartheta\right) e^{im\varphi},\ \varphi \in \left[0, 2\pi\right],\ \vartheta \in \left[0, \pi\right] \tag{24}$$

$$Y_\ell^m\left(\vartheta,\varphi\right)^* = \left(-1\right)^m Y_\ell^{-m}\left(\vartheta,\varphi\right).$$

In particular $Y_\ell^0\left(\vartheta,\varphi\right) = \sqrt{\frac{2\ell+1}{4\pi}} P_\ell\left(\cos\vartheta\right)$, $Y_0^0\left(\vartheta,\varphi\right) = \sqrt{\frac{1}{4\pi}}$,

$$Y_\ell^m\left(N\right) = \delta_{m,0}\sqrt{\frac{2\ell+1}{4\pi}},$$

we have (37)

$$\int_0^{2\pi}\int_0^{\pi} \left|Y_\ell^m\left(\vartheta,\varphi\right)\right|^2 \sin\vartheta d\vartheta d\varphi = 1$$

note that Y_ℓ^m is normalized *fully*, some authors do not apply $1/\sqrt{4\pi}$ in the definition of Y_ℓ^m, also for a sphere with radius R *spherical harmonics are normalized additionally* $Y_\ell^m\left(\vartheta,\varphi\right)/R$.

12. **Wigner $3j$-symbols**, [Lou06] notation

$$\begin{pmatrix} \ell_{1:3} \\ m_{1:3} \end{pmatrix} = \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & m_3 \end{pmatrix},$$

Clebsch-Gordan coefficients,

$$\begin{aligned} C^{\ell,k}_{\ell_1,k_1;\ell_2,k_2} &= (-1)^{\ell_1-\ell_2+k}\sqrt{2\ell+1}\begin{pmatrix} \ell_{1:2} & \ell \\ k_{1:2} & -k \end{pmatrix} \\ \begin{pmatrix} \ell_{1:3} \\ m_{1:3} \end{pmatrix} &= \frac{(-1)^{\ell_3+m_3+2\ell_1}}{\sqrt{2\ell_3+1}} C^{\ell_3,k_3}_{\ell_1,-k_1;\ell_2,-k_2} \end{aligned}$$

$$\begin{aligned} \begin{pmatrix} \ell_{1:3} \\ m_{1:3} \end{pmatrix} &= (-1)^{\ell_1+\ell_2+\ell_3}\begin{pmatrix} \ell_2 & \ell_1 & \ell_3 \\ m_2 & m_1 & m_3 \end{pmatrix} \\ &= \begin{pmatrix} \ell_2 & \ell_3 & \ell_1 \\ m_2 & m_3 & m_1 \end{pmatrix} \\ \begin{pmatrix} \ell_{1:3} \\ m_{1:3} \end{pmatrix} &= (-1)^{\ell_1+\ell_2+\ell_3}\begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ -m_1 & -m_2 & -m_3 \end{pmatrix} \end{aligned} \tag{25}$$

Orthogonality relations

$$\sum_{m_{1:3}} \begin{pmatrix} \ell_{1:3} \\ m_{1:3} \end{pmatrix}^2 = 1 \tag{26}$$

$$(2\ell+1)\sum_{m_{1:2}} \begin{pmatrix} \ell_{1:2} & \ell \\ m_{1:2} & m \end{pmatrix}\begin{pmatrix} \ell_{1:2} & j \\ m_{1:2} & k \end{pmatrix} = \delta_{m,k}\delta_{\ell,j} \tag{27}$$

$$\sum_{\ell,k}(2\ell+1)\begin{pmatrix} \ell_{1:2} & \ell \\ m_{1:2} & k \end{pmatrix}\begin{pmatrix} \ell_{1:2} & \ell \\ k_{1:2} & k \end{pmatrix} = \delta_{m_1,k_1}\delta_{m_2,k_2} \tag{28}$$

$$\begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ 0 & 0 & 0 \end{pmatrix} = 0,$$

if $\mathcal{L} = \ell_1+\ell_2+\ell_3$ is odd, otherwise see [Edm57], (3.7.17)

$$\begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ 0 & 0 & 0 \end{pmatrix} = (-1)^{\mathcal{L}/2}\sqrt{\frac{\prod(\mathcal{L}-2\ell_j)!}{(\mathcal{L}+1)!}}\frac{(\mathcal{L}/2)!}{\prod(\mathcal{L}/2-\ell_j)!},$$

if ℓ is even let $\ell = 2k$, say then

$$\begin{pmatrix} \ell & \ell & \ell \\ 0 & 0 & 0 \end{pmatrix} = (-1)^{3k}\sqrt{\frac{(\ell!)^3}{(6k+1)!}}\frac{(3k)!}{(k!)^3}.$$

13. **Selection rules:**
    - $3j$-symbol **vanishes** unless $m_1 + m_2 + m_3 = 0$.
    - Integer perimeter rule $\ell_1 + \ell_2 + \ell_3$ is an integer (an even if $m_1 = m_2 = m_3 = 0$)
    - triangular inequality $|\ell_1 - \ell_2| \leq \ell_3 \leq \ell_1 + \ell_2$

14. **Gaunt series** [Edm57], (4.6.5)

$$\begin{aligned} Y_{\ell_1}^{m_1}(\vartheta,\varphi)\, Y_{\ell_2}^{m_2}(\vartheta,\varphi) &= \sum_{\ell,m} \sqrt{\frac{(2\ell_1+1)(2\ell_2+1)(2\ell+1)}{4\pi}} \\ &\quad\times \begin{pmatrix} \ell_1 & \ell_2 & \ell \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} \ell_{1:2} & \ell \\ m_{1:2} & m \end{pmatrix} Y_\ell^{m*}(\vartheta,\varphi), \\ Y_{\ell_1}^{m_1}(\vartheta,\varphi)\, Y_{\ell_2}^{m_2}(\vartheta,\varphi) &= \sum_{\ell,m} \sqrt{\frac{(2\ell_1+1)(2\ell_2+1)}{4\pi(2\ell+1)}} C^{\ell,m}_{\ell_1,m_1;\ell_2,m_2} C^{\ell_3,0}_{\ell_1,0;\ell_2,0} Y_\ell^m(\vartheta,\varphi) \end{aligned}$$

    [VMK88], pp. 144, products of three and more

15. **Rayleigh plane wave expansion in 3D:** $\widehat{\underline{k}} = \underline{k}/|\underline{k}|$, $\widehat{\underline{x}} = \underline{x}/|\underline{x}|$, $r = |\underline{x}|$, $k = |\underline{k}|$,

$$e^{i\underline{k}\cdot\underline{x}} = e^{ikr\cos\vartheta} = 4\pi \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} i^\ell j_\ell(kr)\, Y_\ell^m\left(\widehat{\underline{k}}\right)^* Y_\ell^m(\widehat{\underline{x}})$$

    where

$$j_\ell(kr) = \sqrt{\frac{\pi}{2kr}} J_{\ell+1/2}(kr),$$

    is the Spherical Bessel function of the first kind, also

$$\begin{aligned} e^{iz\cos\vartheta} &= \sum_{\ell=-\infty}^{\infty} i^\ell J_\ell(z)\, e^{i\ell\vartheta}, \\ e^{iz\cos\vartheta} &= J_0(z) + \sum_{\ell=-\infty}^{\infty} i^\ell J_\ell(z) \cos\ell\vartheta, \end{aligned}$$

    see [Lou06].

16. **2-D Dirac delta on sphere**

$$\delta(L_1, L_2) = \sum_{\ell,m} Y_\ell^m(L_1)\, Y_\ell^m(L_2) = \frac{1}{4\pi} \sum_\ell (2\ell+1)\, P_\ell(L_1 \cdot L_2),$$

    1-D Dirac delta

$$\delta(x-y) = \frac{1}{2} \sum_\ell (2\ell+1)\, P_\ell(x)\, P_\ell(y),$$

    see [Lou06].

17. **Rotational invariant functions.** The function

$$\begin{aligned} I_{\ell}\left(L_{1},L_{2}\right) &= \frac{4\pi}{2\ell+1}\sum_{k=-\ell}^{\ell}\left(-1\right)^{k}Y_{\ell}^{k}\left(L_{1}\right)Y_{\ell}^{-k}\left(L_{2}\right)\\ &= \frac{4\pi}{2\ell+1}\sum_{k=-\ell}^{\ell}Y_{\ell}^{k}\left(L_{1}\right)Y_{\ell}^{k*}\left(L_{2}\right) \end{aligned} \tag{29}$$

is rotational invariant. This expression is valid when we apply rotation g_{L_1}, where $.g_{L_1}L_1 = N$

$$\begin{aligned} I_{\ell}\left(L_{1},L_{2}\right) &= \frac{4\pi}{2\ell+1}\frac{2\ell+1}{4\pi}\sum_{k=-\ell}^{\ell}\delta_{k,0}Y_{\ell}^{k}\left(g_{L_1}L_{2}\right)\\ &= Y_{\ell}^{0}\left(g_{L_1}L_{2}\right)\\ &= P_{\ell}\left(\cos L_{1}\cdot L_{2}\right). \end{aligned}$$

It follows from the addition formula (39) as well. The function

$$I_{\ell_1\ell_2\ell_3}\left(L_1,L_2,L_3\right) = \frac{\left(4\pi\right)^{3/2}}{\sqrt{\prod\left(2\ell_j+1\right)}}\sum_{m_{1:3}}\begin{pmatrix}\ell_{1:3}\\ m_{1:3}\end{pmatrix}Y_{\ell_1}^{m_1}\left(L_1\right)Y_{\ell_2}^{m_2}\left(L_2\right)Y_{\ell_3}^{m_3}\left(L_3\right) \tag{30}$$

of three location is rotational invariant, see [Lou06], pp.14. We repeat the notation $\begin{pmatrix}\ell_{1:3}\\ m_{1:3}\end{pmatrix} = \begin{pmatrix}\ell_1 & \ell_2 & \ell_3\\ m_1 & m_2 & m_3\end{pmatrix}$. In particular we have

$$I_{\ell_{1:3}}\left(L_{1:3}\right) = I_{\ell_{1:3}}\left(N, g_{L_1}L_2, g_{L_1}L_3\right).$$

18. **Wigner D-matrix** $\Lambda\left(g\right)Y_{\ell}^{m}\left(L\right) = Y_{\ell}^{m}\left(g^{-1}L\right)$

$$\Lambda\left(g\right)Y_{\ell}^{m}\left(L\right) = \sum_{k=-\ell}^{\ell}D_{k,m}^{(\ell)}\left(g\right)Y_{\ell}^{k}\left(L\right) \tag{31}$$

if ℓ is fixed $D_{m,k}^{(\ell)}\left(g\right)$ is unitary

$$\begin{aligned} \sum_{k=-\ell}^{\ell}D_{m_1,k}^{(\ell)}\left(g\right)D_{m_2,k}^{(\ell)*}\left(g\right) &= \delta_{m_1,m_2},\\ \sum_{k=-\ell}^{\ell}D_{k,m_1}^{(\ell)}\left(g\right)D_{k,m_2}^{(\ell)*}\left(g\right) &= \delta_{m_1,m_2},\\ \sum_{k=-\ell}^{\ell}D_{k,m_1}^{(\ell)*}\left(g\right)D_{k,m_2}^{(\ell)}\left(g\right) &= \delta_{m_1,m_2}, \end{aligned}$$

$\left[D_{m,k}^{(\ell)}\left(g\right)\right]^{*}=\left[D_{m,k}^{(\ell)}\left(g\right)\right]^{-1}$. We introduce the notation $D^{(\ell)}=\left[D_{m,k}^{(\ell)}\right]$, for fixed rotation g. Thus $D^{(\ell)}$ denotes a unitary matrix of order $2\ell+1$, and it follows $D^{(\ell)}\left[D^{(\ell)}\right]^{-1}=D^{(\ell)}D^{(\ell)*}$, unimodular $\det D^{(\ell)}=1$. For the definition of $D_{m,k}^{(\ell)}\left(g\right)$

$$D_{m,k}^{(\ell)}\left(\varphi,\vartheta,\gamma\right)=\exp\left(-im\varphi\right)d_{m,k}^{(\ell)}\left(\vartheta\right)\exp\left(-ik\gamma\right) \tag{32}$$

where $d_{m,k}^{(\ell)}$ is the Wigner (small) d-matrix it is real and

$$\begin{aligned}
d_{m,k}^{(\ell)}\left(-\vartheta\right) &= \left(-1\right)^{m-k}d_{m,k}^{(\ell)}\left(\vartheta\right)\\
&= d_{k,m}^{(\ell)}\left(\vartheta\right)\\
&= \left(-1\right)^{k-m}d_{-k,-m}^{(\ell)}\left(\vartheta\right)\\
&= d_{-m,-k}^{(\ell)}\left(\vartheta\right)
\end{aligned}$$

see [VMK88], pp79 for details
The notation $*$ defined as the transpose and conjugate for a matrix and just conjugate for a scalar, for instance

$$\begin{aligned}
\left[D_{m,k}^{(\ell)}\left(\varphi,\vartheta,\gamma\right)\right]^{*} &= \left[\overline{D_{k,m}^{(\ell)}}\left(\varphi,\vartheta,\gamma\right)\right]=\left[\left(-1\right)^{m-k}\overline{D_{m,k}^{(\ell)}}\left(\gamma,\vartheta,\varphi\right)\right]\\
&= \left[\left(-1\right)^{m-k}D_{-m,-k}^{(\ell)}\left(\gamma,\vartheta,\varphi\right)\right]\\
&= \left[D_{m,k}^{(\ell)}\left(-\gamma,-\vartheta,-\varphi\right)\right],
\end{aligned}$$

$$\begin{aligned}
D_{k,m}^{(\ell)}\left(\varphi,\vartheta,\gamma\right) &= \left(-1\right)^{m-k}D_{m,k}^{(\ell)}\left(\gamma,\vartheta,\varphi\right)\\
&= D_{m,k}^{(\ell)}\left(\gamma,-\vartheta,\varphi\right)
\end{aligned}$$

note here that if the Euler angles $(\varphi,\vartheta,\gamma)$ corresponds to g then the one $(-\gamma,-\vartheta,-\varphi)$ corresponds to g^{-1}, hence

$$\begin{aligned}
D_{m,k}^{(\ell)}\left(g\right) &= \left(-1\right)^{m-k}D_{-m,-k}^{(\ell)*}\left(g\right)\\
D_{m,k}^{(\ell)*}\left(g\right) &= D_{k,m}^{(\ell)}\left(g^{-1}\right)
\end{aligned}$$

also

$$D_{m,k}^{(\ell)}\left(g\right)=D_{\ell-m+1,\ell-k+1}^{(\ell)}\left(g\right)$$

Let g the successive application of g_1 and g_2 then

$$D_{m,k}^{(\ell)}\left(g\right)=\sum_{j=-\ell}^{\ell}D_{m,j}^{(\ell)}\left(g_1\right)D_{j,k}^{(\ell)}\left(g_2\right).$$

Condon and Shortley phase convention, [Edm57], (4.3.3)

$$\begin{aligned} Y_\ell^m(\vartheta,\varphi) &= \sqrt{\frac{2\ell+1}{4\pi}} D^{(\ell)}_{0,-m}(\gamma,\vartheta,\varphi) \\ &= (-1)^m D^{(\ell)*}_{0,m}(\gamma,\vartheta,\varphi) \\ &= (-1)^m \sqrt{\frac{2\ell+1}{4\pi}} D^{(\ell)}_{-m,0}(\varphi,\vartheta,\gamma) \\ &= \sqrt{\frac{2\ell+1}{4\pi}} D^{(\ell)*}_{m,0}(\varphi,\vartheta,\gamma), \end{aligned} \tag{33}$$

where γ is arbitrary angle. This form is referred to as passive convention as well. It also follows

$$\begin{aligned} Y_\ell^{m*}(\vartheta,\varphi) &= Y_\ell^m(\vartheta,-\varphi) \\ &= (-1)^m Y_\ell^{-m}(\vartheta,\varphi), \\ Y_\ell^{-m}(\vartheta,\varphi) &= (-1)^m e^{-i2m\varphi} Y_\ell^m(\vartheta,\varphi). \end{aligned}$$

[Wor], (52) in terms of Wigner $3j$-symbols, [Edm57] (4.3.2) Singly coupled form, Clebsch-Gordan series:

$$\begin{aligned} D^{(\ell_1)}_{m_1,k_1} D^{(\ell_2)}_{m_2,k_2} &= \sum_{\ell,m,k} (2\ell+1) \begin{pmatrix} \ell_{1:2} & \ell \\ m_{1:2} & m \end{pmatrix} D^{(\ell)*}_{m,k} \begin{pmatrix} \ell_{1:2} & \ell \\ k_{1:2} & k \end{pmatrix} \\ &= \sum_{\ell,m,k} C^{\ell,m}_{\ell_1,m_1;\ell_2,m_2} C^{\ell,k}_{\ell_1,k_1;\ell_2,k_2} D^{(\ell)}_{m,k}, \end{aligned} \tag{34}$$

$$D^{(\ell_1)}_{m_1,k_1} D^{(\ell_2)}_{m_2,k_2} = \sum_{\ell=|\ell_1-\ell_2|}^{\ell_1+\ell_2} \begin{pmatrix} \ell_{1:2} & \ell \\ m_{1:2} & m \end{pmatrix} \begin{pmatrix} \ell_{1:2} & \ell \\ k_{1:2} & k \end{pmatrix} (-1)^{m-k} (2\ell+1) D^{(\ell)}_{m_1+m_2,k_1+k_2}$$

$-m = m_1 + m_2, -k = k_1 + k_2$, [Edm57], (4.3.4). Double coupled form

$$\sum_{m_1,m_2,k_1,k_2} \begin{pmatrix} \ell_{1:2} & \ell \\ m_{1:2} & m \end{pmatrix} D^{(\ell_1)}_{m_1,k_1} D^{(\ell_2)}_{m_2,k_2} \begin{pmatrix} \ell_{1:2} & j \\ k_{1:2} & k \end{pmatrix} = \frac{\delta_{\ell,j}}{2\ell+1} D^{(\ell)*}_{m,k}$$

[Edm57], (4.3.3)

$$\sum_{m_1,m_2,m_3} D^{(\ell_1)}_{m_1,k_1} D^{(\ell_2)}_{m_2,k_2} D^{(\ell_3)}_{m_3,k_3} \begin{pmatrix} \ell_{1:3} \\ m_{1:3} \end{pmatrix} = \begin{pmatrix} \ell_{1:3} \\ k_{1:3} \end{pmatrix}, \tag{35}$$

The integrals $dg = \sin\vartheta d\vartheta d\varphi d\gamma/8\pi^2$

$$\int_{SO(3)} D_{m,k}^{(\ell)} dg = \delta_{\ell,0}\delta_{m,0}\delta_{k,0} \tag{36}$$

$$\mathcal{G}_{k_1,k_2;m_1,m_2,}^{\ell_1,\ell_2} = \int_{SO(3)} D_{m_1,k_1}^{(\ell_1)*} D_{m_2,k_2}^{(\ell_2)} dg = \delta_{\ell_1,\ell_2}\delta_{m_1,m_2}\delta_{k_1,k_2}\frac{1}{2\ell_1+1} \tag{37}$$

$$\mathcal{G}_{k_1,k_2,k_3;m_1,m_2,m_3}^{\ell_1,\ell_2,\ell_3} = \int_{SO(3)} D_{m_1,k_1}^{(\ell_1)} D_{m_2,k_2}^{(\ell_2)} D_{m_3,k_3}^{(\ell_3)} dg = \begin{pmatrix} \ell_{1:3} \\ m_{1:3} \end{pmatrix} \begin{pmatrix} \ell_{1:3} \\ k_{1:3} \end{pmatrix} \tag{38}$$

$$\int_{SO(3)} D_{m_1,k_1}^{(\ell_1)*} D_{m_2,k_2}^{(\ell_2)} D_{m_3,k_3}^{(\ell_3)} dg = \frac{1}{2\ell_3+1} C_{\ell_1,m_1;\ell_2,m_2}^{\ell_3,m_3} C_{\ell_1,k_1;\ell_2,k_2}^{\ell_3,k_3}$$

where the Haar measure is $dg = \sin\vartheta d\vartheta d\varphi d\gamma/8\pi^2$,[VMK88].

19. Scalar product for f, g

$$(f,g) = \int_0^{2\pi}\int_0^{\pi} f(\vartheta,\varphi)\overline{g(\vartheta,\varphi)}\sin\vartheta d\vartheta d\varphi$$

20. **Addition formula** [GR00]GR 8.814,[EMOT81] vol. 2, 11.4(8) pp. 236, [Ml66]. .$P_\ell(1) = 1$, $\cos\vartheta = ((\vartheta_1,\varphi_1)\cdot(\vartheta_2,\varphi_2))$;

$$\sum_{m=-\ell}^{\ell} Y_\ell^{m*}(\vartheta_1,\varphi_1) Y_\ell^m(\vartheta_2,\varphi_2) = \frac{2\ell+1}{4\pi}P_\ell(\cos\vartheta) \tag{39}$$

[Yad83] For general dimension d [Yad83] p.72 provides an addition formula in terms of Gegenbauer polynomials C_ℓ^ν, $\left(C_\ell^{1/2} = P_\ell\right)$.

C Proofs, Bispectrum

We use the following notations $Z_{\ell_{1:3}}^{m_{1:3}} = \left(Z_{\ell_1}^{m_1}, Z_{\ell_2}^{m_2}, Z_{\ell_3}^{m_3}\right)$, $\begin{pmatrix} \ell_{1:3} \\ m_{1:3} \end{pmatrix} = \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & m_3 \end{pmatrix}$, $B_3(\ell_{1:3}) = B_3(\ell_1,\ell_2,\ell_3)$.

Proof of Lemma. 14

$$\mathrm{Cum}_3\left(Z_{\ell_{1:3}}^{m_{1:3}}\right) = \begin{pmatrix} \ell_{1:3} \\ m_{1:3} \end{pmatrix} B_3(\ell_{1:3}),$$

Assume then

$$\begin{aligned}
\operatorname{Cum}_3\left(\mathcal{Z}_{\ell_{1:3}}^{k_{1:3}}\right) &= \sum_{m_1,m_2,m_3} D_{k_1,m_1}^{(\ell_1)} D_{k_2,m_2}^{(\ell_2)} D_{m_3,k_3}^{(\ell_3)} \operatorname{Cum}_3\left(Z_{\ell_{1:3}}^{m_{1:3}}\right) \qquad (40) \\
&= \sum_{m_{1:3}} D_{k_1,m_1}^{(\ell_1)} D_{k_2,m_2}^{(\ell_2)} D_{m_3,k_3}^{(\ell_3)} \binom{\ell_{1:3}}{m_{1:3}} B_3\left(\ell_{1:3}\right) \\
&= \binom{\ell_{1:3}}{k_{1:3}} B_3\left(\ell_{1:3}\right) \\
&= \operatorname{Cum}_3\left(Z_{\ell_1}^{k_1}, Z_{\ell_2}^{k_2}, Z_{\ell_3}^{k_3}\right)
\end{aligned}$$

$\mathcal{Z}_{\ell_{1:3}}^{k_{1:3}} = \left(\mathcal{Z}_{\ell_1}^{k_1}, \mathcal{Z}_{\ell_2}^{k_2}, \mathcal{Z}_{\ell_3}^{k_3}\right)$ see Edmonds (4.3.3)!![Edm57], [CM01], hence the assumption of isotropy fulfils. If (9) is not assumed then under the assumption of isotropy we have $\operatorname{Cum}_3\left(\mathcal{Z}_{\ell_{1:3}}^{k_{1:3}}\right) = \operatorname{Cum}_3\left(Z_{\ell_{1:3}}^{k_{1:3}}\right)$, and integrate both sides of (40), see (38),

$$\begin{aligned}
\operatorname{Cum}_3\left(Z_{\ell_{1:3}}^{k_{1:3}}\right) &= \binom{\ell_{1:3}}{k_{1:3}} \sum_{m_{1:3}} \binom{\ell_{1:3}}{m_{1:3}} \operatorname{Cum}_3\left(Z_{\ell_{1:3}}^{m_{1:3}}\right) \\
&= \binom{\ell_{1:3}}{k_{1:3}} B_3\left(\ell_{1:3}\right)
\end{aligned}$$

where

$$B_3\left(\ell_{1:3}\right) = \sum_{m_{1:3}} \binom{\ell_{1:3}}{m_{1:3}} \operatorname{Cum}_3\left(Z_{\ell_{1:3}}^{m_{1:3}}\right).$$

Hence (9) is a necessary and sufficient assumption for the third order isotropy. In this case the bispectrum is a linear combination of the cumulants fo the angular projections by the probability amplitude of coupling three angular momenta $\ell_{1:3}$. ■

Proof. Consider the expression

$$\binom{\ell_{1:3}}{k_{1:3}} B_3\left(\ell_{1:3}\right) = \binom{\ell_{1:3}}{k_{1:3}} \sum_{m_{1:3}} \binom{\ell_{1:3}}{m_{1:3}} \operatorname{Cum}_3\left(Z_{\ell_{1:3}}^{m_{1:3}}\right).$$

Any permutation of $\ell_{1:3}$ implies the same sign of the Wigner coefficients hence it does not change the value. ■

Proof. 3-product of spherical harmonics is **rotation invariant,** indeed

$$\begin{aligned}
I_{\ell_{1:3}}\left(gL_{1:3}\right) &= \sum_{m_{1:3}} \binom{\ell_{1:3}}{m_{1:3}} \sum_{k_{1:3}} D_{k_1,m_1}^{(\ell_1)} D_{k_2,m_2}^{(\ell_2)} D_{m_3,k_3}^{(\ell_3)} Y_{\ell_1}^{k_1}\left(L_1\right) Y_{\ell_2}^{k_2}\left(L_2\right) Y_{\ell_3}^{k_3}\left(L_3\right) \\
&= \sum_{k_{1:3}} \binom{\ell_{1:3}}{k_{1:3}} Y_{\ell_1}^{k_1}\left(L_1\right) Y_{\ell_2}^{k_2}\left(L_2\right) Y_{\ell_3}^{k_3}\left(L_3\right).
\end{aligned}$$

see (35). ■

D Proofs, Trispectrum

Repeat the notations $\ell_{1:4}=(\ell_1,\ell_2,\ell_3,\ell_4)$,
$\operatorname{Cum}_4\left(X\left(L_1\right),X\left(L_2\right),X\left(L_3\right),X\left(L_4\right)\right)=\operatorname{Cum}_4\left(X\left(L_{1:4}\right)\right)$, $T_4\left(\ell_1,\ell_2,\ell_3,\ell_4|\,\ell^1\right)=T_4\left(\ell_{1:4}|\,\ell^1\right)$.

Lemma 25

$$\sum_{m_{1:4}}D_{k_1,m_1}^{(\ell_1)}D_{k_2,m_2}^{(\ell_2)}D_{k_3,m_3}^{(\ell_3)}D_{k_4,m_4}^{(\ell_4)}\left(-1\right)^m\begin{pmatrix}\ell_{1:2} & \ell\\ m_{1:2} & -m\end{pmatrix}\begin{pmatrix}\ell & \ell_{3:4}\\ m & m_{3:4}\end{pmatrix}$$
$$=\left(-1\right)^k\begin{pmatrix}\ell_{1:2} & \ell\\ k_{1:2} & -k\end{pmatrix}\begin{pmatrix}\ell_{3:4} & \ell\\ k_{3:4} & k\end{pmatrix}.$$

Proof. Indeed, from

$$D_{k_3,m_3}^{(\ell_3)}D_{k_4,m_4}^{(\ell_4)}=\sum_{\ell^2,k^2,m^2}\left(2\ell^2+1\right)\begin{pmatrix}\ell_{3:4} & \ell^2\\ m_{3:4} & m^2\end{pmatrix}\begin{pmatrix}\ell_{3:4} & \ell^2\\ k_{3:4} & k^2\end{pmatrix}D_{k^2,m^2}^{(\ell^2)*},$$

and from $m_1+m_2+m_3+m_4=0$,follows

$$\begin{aligned}&\sum_{m_{1:4}}D_{k_1,m_1}^{(\ell_1)}D_{k_2,m_2}^{(\ell_2)}D_{k_3,m_3}^{(\ell_3)}D_{k_4,m_4}^{(\ell_4)}\left(-1\right)^m\begin{pmatrix}\ell_{1:2} & \ell\\ m_{1:2} & -m\end{pmatrix}\begin{pmatrix}\ell & \ell_{3:4}\\ m & m_{3:4}\end{pmatrix}\\
&=\sum_{\ell^2,k^2,m^2}\sum_{m_{1:4}}D_{k_1,m_1}^{(\ell_1)}D_{k_2,m_2}^{(\ell_2)}D_{k^2,m^2}^{(\ell^2)*}\left(2\ell^2+1\right)\\
&\times\left(-1\right)^m\begin{pmatrix}\ell_{3:4} & \ell^2\\ m_{3:4} & m^2\end{pmatrix}\begin{pmatrix}\ell_{3:4} & \ell^2\\ k_{3:4} & k^2\end{pmatrix}\begin{pmatrix}\ell_{1:2} & \ell\\ m_{1:2} & -m\end{pmatrix}\begin{pmatrix}\ell & \ell_{3:4}\\ m & m_{3:4}\end{pmatrix}\\
&=\sum_{\ell^2,k^2,m^2}\sum_{m_{3:4}}\begin{pmatrix}\ell_{3:4} & \ell^2\\ m_{3:4} & -m^2\end{pmatrix}\begin{pmatrix}\ell & \ell_{3:4}\\ m & m_{3:4}\end{pmatrix}\left(2\ell^2+1\right)\\
&\times\left(-1\right)^m\sum_{m_{1:2}}D_{k_1,m_1}^{(\ell_1)}D_{k_2,m_2}^{(\ell_2)}D_{k^2,m^2}^{(\ell^2)}\left(-1\right)^{m^2-k^2}\begin{pmatrix}\ell_{3:4} & \ell^2\\ k_{3:4} & -k^2\end{pmatrix}\begin{pmatrix}\ell_{1:2} & \ell\\ m_{1:2} & -m\end{pmatrix}\\
&=\delta_{\ell,\ell^2}\delta_{m,-m^2}\sum_{\ell^2,k^2}\sum_{m^2,m_{1:2}}D_{k_1,m_1}^{(\ell_1)}D_{k_2,m_2}^{(\ell_2)}D_{-k^2,m^2}^{(\ell^2)}\begin{pmatrix}\ell_{1:2} & \ell^2\\ m_{1:2} & -m^2\end{pmatrix}\left(-1\right)^{-k^2}\begin{pmatrix}\ell_{3:4} & \ell^2\\ k_{3:4} & -k^2\end{pmatrix}\\
&=\begin{pmatrix}\ell_{1:2} & \ell\\ k_{1:2} & k^2\end{pmatrix}\left(-1\right)^{-k^2}\begin{pmatrix}\ell_{3:4} & \ell\\ k_{3:4} & -k^2\end{pmatrix}.\end{aligned}$$

Now use (35)

$$\left(2\ell+1\right)\sum_{m_{1:2}}\begin{pmatrix}\ell_{1:2} & \ell\\ m_{1:2} & -m\end{pmatrix}\begin{pmatrix}\ell_{1:2} & \ell^1\\ m_{1:2} & m^1\end{pmatrix}=\delta_{\ell,\ell^1}\delta_{m,-m^1},$$

see (27). ■

Lemma 26

$$\begin{aligned}&\int_{SO(3)} D^{(\ell_1)}_{k_1,m_1} D^{(\ell_2)}_{k_2,m_2} D^{(\ell_3)}_{k_3,m_3} D^{(\ell_4)}_{k_4,m_4} dg \\ &= \sum_{\ell} (2\ell+1)(-1)^{k-m} \begin{pmatrix}\ell_{1:2} & \ell \\ k_{1:2} & -k\end{pmatrix} \begin{pmatrix}\ell_{3:4} & \ell \\ k_{3:4} & k\end{pmatrix} \begin{pmatrix}\ell_{1:2} & \ell \\ m_{1:2} & -m\end{pmatrix} \begin{pmatrix}\ell_{3:4} & \ell \\ m_{3:4} & m\end{pmatrix}\end{aligned}$$

Proof. Here the we apply the formula (34) for the single coupled D-matrices then the integral of triple product of D-matrices 38 and get

$$\begin{aligned}&\int_{SO(3)} D^{(\ell_1)}_{k_1,m_1} D^{(\ell_2)}_{k_2,m_2} D^{(\ell_3)}_{k_3,m_3} D^{(\ell_4)}_{k_4,m_4} dg \\ &= \sum_{\ell^1,k^1,m^1} \left(2\ell^1+1\right)(-1)^{m^1-k^1} \begin{pmatrix}\ell_{3:4} & \ell^1 \\ k_{3:4} & k^1\end{pmatrix} \begin{pmatrix}\ell_{3:4} & \ell^1 \\ m_{3:4} & m^1\end{pmatrix} \int_{SO(3)} D^{(\ell_1)}_{k_1,m_1} D^{(\ell_2)}_{k_2,m_2} D^{(\ell^1)}_{-k^1,-m^1} dg \\ &= \sum_{\ell^1,k^1,m^1} \begin{pmatrix}\ell_{1:2} & \ell^1 \\ k_{1:2} & k^1\end{pmatrix} \begin{pmatrix}\ell_{3:4} & \ell^1 \\ k_{3:4} & -k^1\end{pmatrix} \\ &\times \left(2\ell^1+1\right)(-1)^{m^1-k^1} \begin{pmatrix}\ell_{1:2} & \ell^1 \\ m_{1:2} & m^1\end{pmatrix} \begin{pmatrix}\ell_{3:4} & \ell^1 \\ m_{3:4} & -m^1\end{pmatrix}\end{aligned}$$

■

Lemma 27 *If* $\ell_1+\ell_2+\ell_3+\ell_4$ *is even then*

$$\sum_{\ell} (2\ell+1)(-1)^{k-m} \begin{pmatrix}\ell_{1:2} & \ell \\ k_{1:2} & -k\end{pmatrix} \begin{pmatrix}\ell & \ell_{3:4} \\ k & k_{3:4}\end{pmatrix} \begin{pmatrix}\ell_{1:2} & \ell \\ m_{1:2} & -m\end{pmatrix} \begin{pmatrix}\ell & \ell_{3:4} \\ m & m_{3:4}\end{pmatrix}$$

is symmetric.

Proof. of Lemma 27. Start with

$$\begin{aligned}&\begin{pmatrix}\ell_{1:2} & \ell \\ k_{1:2} & -k\end{pmatrix} \begin{pmatrix}\ell & \ell_{3:4} \\ k & k_{3:4}\end{pmatrix} \\ &\qquad = \sum_{\ell_0} (2\ell_0+1)(-1)^{\Sigma+\ell_0-\ell-k_1-k_3} \begin{pmatrix}\ell_2 & \ell_3 & \ell_0 \\ k_2 & k_3 & -k_0\end{pmatrix} \begin{pmatrix}\ell_1 & \ell_0 & \ell_4 \\ k_1 & k_0 & k_4\end{pmatrix} \begin{Bmatrix}\ell_1 & \ell_2 & \ell \\ \ell_3 & \ell_4 & \ell_0\end{Bmatrix}\end{aligned}$$

where $\Sigma = \ell_1+\ell_2+\ell_3+\ell_4$, . Now we evaluate the product

$$\begin{aligned}&\sum_{\ell_0^1} \left(2\ell_0^1+1\right)(-1)^{\Sigma+\ell_0^1-\ell-k_1-k_3} \begin{pmatrix}\ell_2 & \ell_3 & \ell_0^1 \\ k_2 & k_3 & -k_0^1\end{pmatrix} \begin{pmatrix}\ell_1 & \ell_0^1 & \ell_4 \\ k_1 & k_0^1 & k_4\end{pmatrix} \begin{Bmatrix}\ell_1 & \ell_2 & \ell \\ \ell_3 & \ell_4 & \ell_0^1\end{Bmatrix} \\ &\times \sum_{\ell_0^2} \left(2\ell_0^2+1\right)(-1)^{\Sigma+\ell_0^2-\ell-m_1-m_3} \begin{pmatrix}\ell_2 & \ell_3 & \ell_0^2 \\ m_2 & m_3 & -m_0^2\end{pmatrix} \begin{pmatrix}\ell_1 & \ell_0^2 & \ell_4 \\ m_1 & m_0^2 & m_4\end{pmatrix} \begin{Bmatrix}\ell_1 & \ell_2 & \ell \\ \ell_3 & \ell_4 & \ell_0^2\end{Bmatrix}.\end{aligned}$$

The values
$$\sqrt{\left(2\ell+1\right)\left(2\ell_0^1+1\right)}\begin{Bmatrix}\ell_1 & \ell_2 & \ell\\ \ell_3 & \ell_4 & \ell_0^1\end{Bmatrix}$$
form a real orthogonal matrix, see [Edm57] (6.2.10), rows and columns being labelled by ℓ and ℓ_0^1. Hence if we sum it up by ℓ, the result is

$$\sum_{\ell_0^1}\left(2\ell_0^1+1\right)(-1)^{-k_0^1}\begin{pmatrix}\ell_2 & \ell_3 & \ell_0^1\\ k_2 & k_3 & -k_0^1\end{pmatrix}\begin{pmatrix}\ell_1 & \ell_0^1 & \ell_4\\ k_1 & k_0^1 & k_4\end{pmatrix}$$
$$\times(-1)^{-m_0^1}\begin{pmatrix}\ell_2 & \ell_3 & \ell_0^1\\ m_2 & m_3 & -m_0^1\end{pmatrix}\begin{pmatrix}\ell_1 & \ell_0^1 & \ell_4\\ m_1 & m_0^1 & m_4\end{pmatrix},$$

q.e.d. ■

Proof of Lemma 16. We have
$$\operatorname{Cum}_4\left(\mathcal{Z}_{\ell_{1:4}}^{k_{1:4}}\right)=\sum_{m_{1:4}}D_{k_1,m_1}^{(\ell_1)}D_{k_2,m_2}^{(\ell_2)}D_{k_3,m_3}^{(\ell_3)}D_{k_4,m_4}^{(\ell_4)}\operatorname{Cum}_4\left(Z_{\ell_{1:4}}^{m_{1:4}}\right).$$

Under the assumption of isotropy $\operatorname{Cum}_4\left(\mathcal{Z}_{\ell_{1:4}}^{k_{1:4}}\right)=\operatorname{Cum}_4\left(Z_{\ell_{1:4}}^{k_{1:4}}\right)$, now integrate both sides by the Haar measure and get

$$\operatorname{Cum}_4\left(Z_{\ell_{1:4}}^{k_{1:4}}\right)=\sum_{m_{1:4}}\sum_{\ell^1,m^1,k^1}\begin{pmatrix}\ell_{1:2} & \ell^1\\ k_{1:2} & -k^1\end{pmatrix}\begin{pmatrix}\ell^1 & \ell_{3:4}\\ k^1 & k_{3:4}\end{pmatrix}(-1)^{k^1-m^1}$$
$$\times\left(2\ell^1+1\right)\begin{pmatrix}\ell_{1:2} & \ell^1\\ m_{1:2} & -m^1\end{pmatrix}\begin{pmatrix}\ell^1 & \ell_{3:4}\\ m^1 & m_{3:4}\end{pmatrix}\operatorname{Cum}_4\left(Z_{\ell_{1:4}}^{m_{1:4}}\right)$$

Note that each term according to summation $m_{1:4}$ is **symmetric** in $\ell_{1:4}$, see Lemma 27 above. Now define
$$T_4\left(\ell_{1:4}|\,\ell^1\right)=\sum_{m^1,m_{1:4}}(-1)^{m^1}\left(2\ell^1+1\right)\begin{pmatrix}\ell_{1:2} & \ell^1\\ m_{1:2} & -m^1\end{pmatrix}\begin{pmatrix}\ell^1 & \ell_{3:4}\\ m^1 & m_{3:4}\end{pmatrix}\operatorname{Cum}_4\left(Z_{\ell_{1:4}}^{m_{1:4}}\right),$$

with this notation we have the cumulant in the form
$$\operatorname{Cum}_4\left(Z_{\ell_{1:4}}^{k_{1:4}}\right)=\sum_{\ell^1,k^1}\begin{pmatrix}\ell_{1:2} & \ell^1\\ k_{1:2} & -k^1\end{pmatrix}\begin{pmatrix}\ell^1 & \ell_{3:4}\\ k^1 & k_{3:4}\end{pmatrix}(-1)^{k^1}T_4\left(\ell_{1:4}|\,\ell^1\right).$$

If instead of isotropy (10) is assumed then

$$\begin{aligned}
\operatorname{Cum}_4\left(\mathcal{Z}^{k_{1:4}}_{\ell_{1:4}}\right) &= \sum_{m_{1:4}} D^{(\ell_1)}_{k_1,m_1} D^{(\ell_2)}_{k_2,m_2} D^{(\ell_3)}_{k_3,m_3} D^{(\ell_4)}_{k_4,m_4} \operatorname{Cum}_4\left(Z^{m_{1:4}}_{\ell_{1:4}}\right) \\
&= \sum_{m_{1:4}} D^{(\ell_1)}_{k_1,m_1} D^{(\ell_2)}_{k_2,m_2} D^{(\ell_3)}_{k_3,m_3} D^{(\ell_4)}_{k_4,m_4} \\
&\times \sum_{\ell} \begin{pmatrix} \ell_{1:2} & \ell \\ m_{1:2} & -m \end{pmatrix} \begin{pmatrix} \ell & \ell_{3:4} \\ m & m_{3:4} \end{pmatrix} (-1)^m (2\ell+1)\, T_4\left(\ell_{1:4}\middle|\,\ell\right) \\
&= \sum_{\ell} T_4\left(\ell_{1:4}\middle|\,\ell\right)(2\ell+1) \\
&\times \sum_{m_{1:4}} D^{(\ell_1)}_{k_1,m_1} D^{(\ell_2)}_{k_2,m_2} D^{(\ell_3)}_{k_3,m_3} D^{(\ell_4)}_{k_4,m_4} (-1)^m \begin{pmatrix} \ell_{1:2} & \ell \\ m_{1:2} & -m \end{pmatrix} \begin{pmatrix} \ell & \ell_{3:4} \\ m & m_{3:4} \end{pmatrix},
\end{aligned}$$

the Lemma 25 can be applied and get

$$\begin{aligned}
\operatorname{Cum}_4\left(\mathcal{Z}^{k_{1:4}}_{\ell_{1:4}}\right) &= \sum_{k,\ell} \begin{pmatrix} \ell_{1:2} & \ell \\ k_{1:2} & -k \end{pmatrix} \begin{pmatrix} \ell_{3:4} & \ell \\ k_{3:4} & k \end{pmatrix} (-1)^k\, T_4\left(\ell_{1:4}\middle|\,\ell^1\right) \\
&= \operatorname{Cum}_4\left(Z^{k_{1:4}}_{\ell_{1:4}}\right).
\end{aligned}$$

■

Particular cases

$$\begin{aligned}
\operatorname{Cum}_4\left(u_{\ell_1}(L), u_{\ell_2}(L), u_{\ell_3}(L), u_{\ell_4}(L)\right) &= \sum_{m^1,\ell^1} T_4\left(\ell_1,\ell_2,\ell_3,\ell_4\middle|\,\ell^1\right) \sum_{m_{1:4}} \prod_{j=1}^{4} Y^{m_j}_{\ell_j}(L_j) \\
&\times \begin{pmatrix} \ell_1 & \ell_2 & \ell^1 \\ m_1 & m_2 & -m^1 \end{pmatrix} \begin{pmatrix} \ell^1 & \ell_3 & \ell_4 \\ m^1 & m_3 & 0 \end{pmatrix},
\end{aligned}$$

$$\begin{aligned}
&\operatorname{Cum}_4\left(u_{\ell_1}(L_1), u_{\ell_2}(L_2), u_{\ell_3}(L), u_{\ell_4}(L)\right) \\
&\quad = \operatorname{Cum}_4\left(u_{\ell_1}(g_{LL_2}L_1), u_{\ell_2}(g_{LL_2}L_2), u_{\ell_3}(N), u_{\ell_4}(N)\right) \\
&= \sqrt{\prod_{j=3:4} \frac{2\ell_j+1}{4\pi}} \sum_{\ell^1} T_4\left(\ell_1,\ell_2,\ell_3,\ell_4\middle|\,\ell^1\right) \begin{pmatrix} \ell_3 & \ell_4 & \ell^1 \\ 0 & 0 & 0 \end{pmatrix} \sqrt{2\ell^1+1} \\
&\quad \times \sum_{m_1,m_2} Y^{m_1}_{\ell_1}(g_{LL_2}L_1)\, Y^{m_2}_{\ell_2}(g_{LL_2}L_2) \begin{pmatrix} \ell_{1:2} & \ell^1 \\ m_{1:2} & 0 \end{pmatrix} \\
&= \frac{\sqrt{2}}{4\pi} \sqrt{\prod_{j=3:4} \frac{2\ell_j+1}{4\pi}} \sum_{\ell^1} \begin{pmatrix} \ell_3 & \ell_4 & \ell^1 \\ 0 & 0 & 0 \end{pmatrix} T_4\left(\ell_1,\ell_2,\ell_3,\ell_4\middle|\,\ell^1\right) \mathcal{I}_{\ell_1,\ell_2,\ell^1}(\vartheta_2,\varphi_2,\vartheta_1),
\end{aligned}$$

$$\begin{aligned}
&\mathrm{Cum}_4\left(u_{\ell_1}\left(L_1\right), u_{\ell_2}\left(L\right), u_{\ell_3}\left(L\right), u_{\ell_4}\left(L\right)\right) = \mathrm{Cum}_4\left(u_{\ell_1}\left(g_{LL_1}L_1\right), u_{\ell_2}\left(N\right), u_{\ell_3}\left(N\right), u_{\ell_4}\left(N\right)\right)\\
&= \sqrt{\prod_{j=2:4}\frac{2\ell_j+1}{4\pi}}\sum_{\ell^1} T_4\left(\ell_1,\ell_2,\ell_3,\ell_4|\,\ell^1\right)\sqrt{2\ell^1+1}Y_{\ell_1}^0\left(g_{LL_1}L_1\right)\begin{pmatrix}\ell_1 & \ell_2 & \ell^1\\ 0 & 0 & 0\end{pmatrix}\begin{pmatrix}\ell^1 & \ell_3 & \ell_4\\ 0 & 0 & 0\end{pmatrix}\\
&= \sqrt{\prod_{j=1:4}\frac{2\ell_j+1}{4\pi}}\sum_{\ell^1} T_4\left(\ell_1,\ell_2,\ell_3,\ell_4|\,\ell^1\right)\sqrt{2\ell^1+1}P_{\ell_1}\left(L\cdot L_1\right)\begin{pmatrix}\ell_1 & \ell_2 & \ell^1\\ 0 & 0 & 0\end{pmatrix}\begin{pmatrix}\ell^1 & \ell_3 & \ell_4\\ 0 & 0 & 0\end{pmatrix},
\end{aligned}$$

$$\begin{aligned}
&\mathrm{Cum}_4\left(u_{\ell_1}\left(L\right), u_{\ell_2}\left(L\right), u_{\ell_3}\left(L\right), u_{\ell_4}\left(L\right)\right) = \mathrm{Cum}_4\left(u_{\ell_1}\left(N\right), u_{\ell_2}\left(N\right), u_{\ell_3}\left(N\right), u_{\ell_4}\left(N\right)\right)\\
&\qquad = \sqrt{\prod_{j=1:4}\frac{2\ell_i+1}{4\pi}}\sum_{\ell^1} T_4\left(\ell_1,\ell_2,\ell_3,\ell_4|\,\ell^1\right)\sqrt{2\ell^1+1}\begin{pmatrix}\ell_1 & \ell_2 & \ell^1\\ 0 & 0 & 0\end{pmatrix}\begin{pmatrix}\ell^1 & \ell_3 & \ell_4\\ 0 & 0 & 0\end{pmatrix}.
\end{aligned}$$

E Proofs, Polyspectrum

Notation: $\mathcal{G}_{k_{1:p},m_{1:p}}^{\ell_{1:p}} = \mathcal{G}_{k_1,k_2,\ldots,k_p,m_1,m_2,\ldots,m_p}^{\ell_1,\ell_2,\ldots,\ell_{3p}}$.

Lemma 28 *Define* $\ell^0=\ell_1$, $\ell^{p-2}=\ell_p$, $k^0=-k_1$, $k^{p-2}=k_p$, $m^0=-m_1$, $m^{p-2}=m_p$

$$\begin{aligned}
\mathcal{G}_{k_{1:p},m_{1:p}}^{\ell_{1:p}} = \int_{SO(3)}\prod_{a=1}^{p} D_{k_a,m_a}^{(\ell_a)}dg = \sum_{\ell^a,m^a,k^a}(-1)^{\Sigma\left(m^{1:p-3}-k^{1:p\infty-3}\right)}\prod_{a=0}^{p-3}\begin{pmatrix}\ell^a & \ell_{a+2} & \ell^{a+1}\\ -k^a & k_{a+2} & k^{a+1}\end{pmatrix}&\\
\times\prod_{a=0}^{p-3}\begin{pmatrix}\ell^a & \ell_{a+2} & \ell^{a+1}\\ -m^a & m_{a+2} & m^{a+1}\end{pmatrix}\prod_{a=1}^{p-3}\left(2\ell^a+1\right),&
\end{aligned}$$

Proof. The fourth order key formula is emphasized here for understanding the induction

$$\begin{aligned}
&\mathcal{G}^{\ell_{1:5}}_{k_{1:5};m_{1:5}} = \int_{SO(3)} D^{(\ell_1)}_{k_1,m_1} D^{(\ell_2)}_{k_2,m_2} D^{(\ell_3)}_{k_3,m_3} D^{(\ell_4)}_{k_4,m_4} D^{(\ell_5)}_{k_5,m_5} dg \\
&= \sum_{\ell^2,k^2,m^2} \int_{SO(3)} D^{(\ell_1)}_{k_1,m_1} D^{(\ell_2)}_{k_2,m_2} D^{(\ell_3)}_{k_3,m_3} D^{(\ell^2)}_{-k^2,-m^2} dg \\
&\times (-1)^{m^2-k^2} \left(2\ell^2+1\right) \begin{pmatrix} \ell_{4:5} & \ell^2 \\ k_{4:5} & k^2 \end{pmatrix} \begin{pmatrix} \ell_{4:5} & \ell^2 \\ m_{4:5} & m^2 \end{pmatrix} \\
&= \sum_{\ell^{1:2},k^{1:2},m^{1:2}} \begin{pmatrix} \ell_{1:2} & \ell^1 \\ k_{1:2} & k^1 \end{pmatrix} \begin{pmatrix} \ell_3 & \ell^2 & \ell^1 \\ k_3 & k^2 & -k^1 \end{pmatrix} \begin{pmatrix} \ell_{4:5} & \ell^2 \\ k_{4:5} & -k^2 \end{pmatrix} (-1)^{m^1-k^1} \\
&\times \left(2\ell^1+1\right) \begin{pmatrix} \ell_{1:2} & \ell^1 \\ m_{1:2} & m^1 \end{pmatrix} \begin{pmatrix} \ell_3 & \ell^2 & \ell^1 \\ m_3 & m^2 & -m^1 \end{pmatrix} \begin{pmatrix} \ell_{4:5} & \ell^2 \\ m_{4:5} & -m^2 \end{pmatrix} \\
&\times (-1)^{m^2-k^2} \left(2\ell^2+1\right) \\
&= \sum_{\ell^{1:2},k^{1:2},m^{1:2}} \begin{pmatrix} \ell_{1:2} & \ell^1 \\ k_{1:2} & k^1 \end{pmatrix} \begin{pmatrix} \ell^1 & \ell_3 & \ell^2 \\ -k^1 & k_3 & k^2 \end{pmatrix} \begin{pmatrix} \ell^2 & \ell_{4:5} \\ -k^2 & k_{4:5} \end{pmatrix} (-1)^{m^1-k^1} \left(2\ell^1+1\right) \\
&\times \begin{pmatrix} \ell_{1:2} & \ell^1 \\ m_{1:2} & m^1 \end{pmatrix} \begin{pmatrix} \ell^1 & \ell_3 & \ell^2 \\ -m^1 & m_3 & m^2 \end{pmatrix} \begin{pmatrix} \ell^2 & \ell_{4:5} \\ -m^2 & m_{4:5} \end{pmatrix} \times (-1)^{m^2-k^2} \left(2\ell^2+1\right).
\end{aligned}$$

Define $\ell^{p-2} = \ell_p$, $k^{p-2} = k_p$, $m^{p-2} = m_p$

$$\int_{SO(3)} \prod_{a=1}^{p} D^{(\ell_a)}_{k_a,m_a} dg = (-1)^{m_p} \sum_{\ell^a,m^a,k^a} \prod_{a=0}^{p-3} C^{\ell^{a+1},k^{a+1}}_{\ell^a,k^a;\ell_{a+2},k_{a+2}} C^{\ell^{a+1},m^{a+1}}_{\ell^a,m^a;\ell_{a+2},m_{a+2}}$$

$$\begin{aligned}
\int_{SO(3)} \prod_{a=1}^{p} D^{(\ell_a)}_{k_a,m_a} dg = \sum_{\ell^a,m^a,k^a} & (-1)^{\Sigma\left(m^{1:p-3}-k^{1:p-3}\right)} \begin{pmatrix} \ell_{1:2} & \ell^1 \\ k_{1:2} & k^1 \end{pmatrix} \prod_{a=1}^{p-3} \begin{pmatrix} \ell^a & \ell_{a+2} & \ell^{a+1} \\ -k^a & k_{a+2} & k^{a+1} \end{pmatrix} \\
& \times \begin{pmatrix} \ell_{1:2} & \ell^1 \\ m_{1:2} & m^1 \end{pmatrix} \prod_{a=1}^{p-3} \begin{pmatrix} \ell^a & \ell_{a+2} & \ell^{a+1} \\ -m^a & m_{a+2} & m^{a+1} \end{pmatrix} \left(2\ell^{a+1}+1\right)
\end{aligned}$$

■

Lemma 29

$$\begin{aligned}
\sum_{m_{1:p}} \prod_{a=1}^{p} D^{(\ell_a)}_{k_a,m_a} & (-1)^{\Sigma m^{1:p-3}} \prod_{a=0}^{p-3} \begin{pmatrix} \ell^a & \ell_{a+2} & \ell^{a+1} \\ -m^a & m_{a+2} & m^{a+1} \end{pmatrix} \\
&= (-1)^{\Sigma\left(m^{1:p-3}-k^{1:p-3}\right)} \prod_{a=0}^{p-3} \begin{pmatrix} \ell^a & \ell_{a+2} & \ell^{a+1} \\ -k^a & k_{a+2} & k^{a+1} \end{pmatrix}.
\end{aligned}$$

Proof. Induction

$$\sum_{m_{1:4}} D^{(\ell_1)}_{k_1,m_1} D^{(\ell_2)}_{k_2,m_2} D^{(\ell_3)}_{k_3,m_3} D^{(\ell_4)}_{k_4,m_4} (-1)^m \begin{pmatrix} \ell_{1:2} & \ell \\ m_{1:2} & -m \end{pmatrix} \begin{pmatrix} \ell & \ell_{3:4} \\ m & m_{3:4} \end{pmatrix} = (-1)^k \begin{pmatrix} \ell_{1:2} & \ell \\ k_{1:2} & -k \end{pmatrix} \begin{pmatrix} \ell_{3:4} & \ell \\ k_{3:4} & k \end{pmatrix}.$$

Indeed

$$D^{(\ell_3)}_{k_3,m_3} D^{(\ell_4)}_{k_4,m_4} = \sum_{\ell^2,k^2,m^2} \left(2\ell^2+1\right) \begin{pmatrix} \ell_{3:4} & \ell^2 \\ m_{3:4} & m^2 \end{pmatrix} \begin{pmatrix} \ell_{3:4} & \ell^2 \\ k_{3:4} & k^2 \end{pmatrix} D^{(\ell^2)*}_{k^2,m^2}$$

$m_1 + m_2 + m_3 + m_4 = 0$

$$\begin{aligned}
&\sum_{m_{1:4}} D^{(\ell_1)}_{k_1,m_1} D^{(\ell_2)}_{k_2,m_2} D^{(\ell_3)}_{k_3,m_3} D^{(\ell_4)}_{k_4,m_4} (-1)^m \begin{pmatrix} \ell_{1:2} & \ell \\ m_{1:2} & -m \end{pmatrix} \begin{pmatrix} \ell & \ell_{3:4} \\ m & m_{3:4} \end{pmatrix} \\
&= \sum_{\ell^2,k^2,m^2} \sum_{m_{1:4}} D^{(\ell_1)}_{k_1,m_1} D^{(\ell_2)}_{k_2,m_2} D^{(\ell^2)*}_{k^2,m^2} \\
&\times \left(2\ell^2+1\right) (-1)^m \begin{pmatrix} \ell_{3:4} & \ell^2 \\ m_{3:4} & m^2 \end{pmatrix} \begin{pmatrix} \ell_{3:4} & \ell^2 \\ k_{3:4} & k^2 \end{pmatrix} \begin{pmatrix} \ell_{1:2} & \ell \\ m_{1:2} & -m \end{pmatrix} \begin{pmatrix} \ell & \ell_{3:4} \\ m & m_{3:4} \end{pmatrix} \\
&= \sum_{\ell^2,k^2,m^2} \sum_{m_{3:4}} \begin{pmatrix} \ell_{3:4} & \ell^2 \\ m_{3:4} & -m^2 \end{pmatrix} \begin{pmatrix} \ell & \ell_{3:4} \\ m & m_{3:4} \end{pmatrix} \left(2\ell^2+1\right) (-1)^m \\
&\times \sum_{m_{1:2}} D^{(\ell_1)}_{k_1,m_1} D^{(\ell_2)}_{k_2,m_2} D^{(\ell^2)}_{k^2,m^2} (-1)^{m^2-k^2} \begin{pmatrix} \ell_{3:4} & \ell^2 \\ k_{3:4} & -k^2 \end{pmatrix} \begin{pmatrix} \ell_{1:2} & \ell \\ m_{1:2} & -m \end{pmatrix} \\
&= \delta_{\ell,\ell^2} \delta_{m,-m^2} \sum_{\ell^2,k^2} \sum_{m^2,m_{1:2}} D^{(\ell_1)}_{k_1,m_1} D^{(\ell_2)}_{k_2,m_2} D^{(\ell^2)}_{-k^2,m^2} \begin{pmatrix} \ell_{1:2} & \ell^2 \\ m_{1:2} & -m^2 \end{pmatrix} (-1)^{-k^2} \begin{pmatrix} \ell_{3:4} & \ell^2 \\ k_{3:4} & -k^2 \end{pmatrix} \\
&= \begin{pmatrix} \ell_{1:2} & \ell \\ k_{1:2} & k^2 \end{pmatrix} (-1)^{-k^2} \begin{pmatrix} \ell_{3:4} & \ell \\ k_{3:4} & -k^2 \end{pmatrix}
\end{aligned}$$

use (35) ■

Proof. of Lemma 19 We have

$$\operatorname{Cum}_p \left(\mathcal{Z}^{k_{1:p}}_{\ell_{1:p}}\right) = \sum_{m_{1:p}} \prod_{a=1}^{p} D^{(\ell_a)}_{k_a,m_a} \operatorname{Cum}_4 \left(Z^{m_{1:p}}_{\ell_{1:p}}\right).$$

Under isotropy assumption

$$
\begin{aligned}
\operatorname{Cum}_p\left(Z^{k_{1:p}}_{\ell_{1:p}}\right) &= \sum_{m_{1:p}} \int_{SO(3)} \prod_{a=1}^{p} D^{(\ell_a)}_{k_a,m_a} dg \operatorname{Cum}_p\left(Z^{m_{1:p}}_{\ell_{1:p}}\right) \\
&= \sum_{\ell^a, m^a, k^a} (-1)^{\Sigma k^{1:p-3}} \prod_{a=0}^{p-3} \begin{pmatrix} \ell^a & \ell_{a+2} & \ell^{a+1} \\ -k^a & k_{a+2} & k^{a+1} \end{pmatrix} \\
&\times \sum_{m_{1:p}} (-1)^{\Sigma m^{1:p-3}} \prod_{a=0}^{p-3} \begin{pmatrix} \ell^a & \ell_{a+2} & \ell^{a+1} \\ -m^a & m_{a+2} & m^{a+1} \end{pmatrix} \left(2\ell^{a+1}+1\right) \operatorname{Cum}_p\left(Z^{m_{1:p}}_{\ell_{1:p}}\right) \\
&= \sum_{\ell^a, m^a, k^a} (-1)^{\Sigma k^{1:p-3}} \prod_{a=0}^{p-3} \begin{pmatrix} \ell^a & \ell_{a+2} & \ell^{a+1} \\ -k^a & k_{a+2} & k^{a+1} \end{pmatrix} C_p\left(\ell_{1:p}\middle| \ell^{1:p-3}\right).
\end{aligned}
$$

If isotropy is not assumed, apply

$$
\begin{aligned}
\operatorname{Cum}_p\left(Z^{m_{1:p}}_{\ell_{1:p}}\right) = &\sum_{\ell^{1:p-3}, m^{1:p-3}} (-1)^{\Sigma m^{1:p-3}} \prod_{a=0}^{p-3} \begin{pmatrix} \ell^a & \ell_{a+2} & \ell^{a+1} \\ -m^a & m_{a+2} & m^{a+1} \end{pmatrix} \\
&\times \prod_{a=1}^{p-3} \sqrt{2\ell^a+1} \widetilde{S}_p\left(\ell_{1:p}\middle| \ell^{1:p-3}\right).
\end{aligned}
$$

We have

$$
\begin{aligned}
\operatorname{Cum}_p\left(\mathcal{Z}^{k_{1:p}}_{\ell_{1:p}}\right) &= \sum_{m_{1:p}} \prod_{a=1}^{p} D^{(\ell_a)}_{k_a,m_a} (-1)^{\Sigma m^{1:p-3}} \prod_{a=0}^{p-3} \begin{pmatrix} \ell^a & \ell_{a+2} & \ell^{a+1} \\ -m^a & m_{a+2} & m^{a+1} \end{pmatrix} \\
&\times \prod_{a=1}^{p-3} \sqrt{2\ell^a+1} \widetilde{S}_p\left(\ell_{1:p}\middle| \ell^{1:p-3}\right) \\
&= (-1)^{\Sigma k^{1:p-3}} \prod_{a=0}^{p-3} \begin{pmatrix} \ell^a & \ell_{a+2} & \ell^{a+1} \\ -k^a & k_{a+2} & k^{a+1} \end{pmatrix} \\
&\times \prod_{a=1}^{p-3} \sqrt{2\ell^a+1} \widetilde{S}_p\left(\ell_{1:p}\middle| \ell^{1:p-3}\right),
\end{aligned}
$$

see Lemma 29 . ■

Acknowledgement 30 *The publication was supported by the TÁMOP-4.2.2.C-11/1/KONV-2012-0001 project. The project has been supported by the European Union, co-financed by the European Social Fund.*

References

[AH12] P. Adshead and W. Hu, *Fast computation of first-order feature-bispectrum corrections*, Phys. Rev. D **85** (2012), 103531.

[Apo07] M. T. Apostol, *Calculus (volume - 2)*, Wiley, 2007.

[AS92] M. Abramowitz and I. A. Stegun, *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, Dover Publications Inc., New York, 1992, Reprint of the 1972 edition.

[Bar99] J.G. Bartlett, *The standard cosmological model and cmb anisotropies*, New Astron.Rev. **43** (1999), 83–109.

[BLSH12] A. Benoit-Lvy, K. M. Smith, and W. Hu, *Non-gaussian structure of the lensed cmb power spectra covariance matrix*, arXiv:1205.0474 [astro-ph.CO] (2012), 16.

[BM07] P. Baldi and D. Marinucci, *Some characterizations of the spherical harmonics coefficients for isotropic random fields*, Statistics & Probability Letters **77** (2007), 490–496.

[Bre63] E. Breitenberger, *Analogues of the normal distribution on the circle and the sphere*, Biometrika **50** (1963), 81–88.

[Bri65] D. R. Brillinger, *An introduction to polyspectra*, Ann. Math. Statistics **36** (1965), 1351–1374.

[Bri01] ______, *Time series; data analysis and theory*, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2001, Reprint of the 1981 edition.

[BSMR10] N. Bartolo, S. S Matarrese, and A. Riotto, *Non-gaussianity and the cosmic microwave background anisotropies*, Advances in Astronomy **2010** (2010), 68.

[CM01] C. R. Contaldi and J. Magueijo, *Generating non-gaussian maps with a given power spectrum and bispectrum*, PHYSICAL REVIEW D **63** (2001), 103512/1–10.

[CW11] N. Cressie and C. K. Wikle, *Statistics for spatio-temporal data*, Wiley Series in Probability and Statistics, 2011.

[Edm57] A. R. Edmonds, *Angular momentum in quantum mechanics*, Princeton University Press, 1957.

[EMOT81] Arthur Erdélyi, Wilhelm Magnus, Fritz Oberhettinger, and Francesco G. Tricomi, *Higher transcendental functions. Vol. I*, Robert E. Krieger Publishing Co. Inc., Melbourne, Fla., 1981, Based on notes left by Harry Bateman, With a preface by Mina Rees, With a foreword by E. C. Watson, Reprint of the 1953 original.

[FS09] J. R. Fergusson and E. P. S. Shellard, *Shape of primordial non-gaussianity and the cmb bispectrum*, Phys. Rev. D **80** (2009), 043510.

[GG10] C. Gaetan and X. Guyon, *Spatial statistics and modeling*, Springer Series in Statistics, vol. 81, Springer Science+Business Media, 2010.

[GR00] I. S. Gradshteyn and I. M. Ryzhik, *Table of integrals, series, and products*, sixth ed., Academic Press Inc., San Diego, CA, 2000, Translated from the Russian, Translation edited and with a preface by Alan Jeffrey and Daniel Zwillinger.

[HD02] Wayne Hu and Scott Dodelson, *Cosmic microwave background anisotropies*, Annual Review of Astronomy and Astrophysics **40** (2002), no. 1, 171–216.

[Hin82] M. J. Hinich, *Testing for Gaussianity and linearity of a stationary time series*, Journal of Time Series Analysis **3** (1982), 169–176, JTSA.

[Hu01] W. Hu, *Angular trispectrum of the cosmic microwave background*, Physical Review D (Particles, Fields, Gravitation, and Cosmology) **64** (2001), 083005.

[IMK12] K. Izumi, S. Mizuno, and K. Koyama, *Trispectrum estimation in various models of equilateral type non-gaussianity*, Phys. Rev. D **85** (2012), 023521.

[Jon63] R. H. Jones, *Stochastic processes on a sphere*, The Annals of Mathematical Statistics **34** (1963), 213–218.

[Jon94] P. D. Jones, *Hemispheric surface air temperature variations: A reanalysis and an update to 1993*, Journal of Climate **7** (1994), 1794–1802.

[Kak12] R. Kakarala, *The bispectrum as a source of phase-sensitive invariants for fourier descriptors: A group-theoretic approach*, Journal of Mathematical Imaging and Vision **44** (2012), 341–353.

[KBID93] R. Kakarala, B.M. Bennett, G.J. Iverson, and M. D'Zmura, *Bispectral techniques for spherical functions*, Acoustics, Speech, and Signal Processing, 1993. ICASSP-93., 1993 IEEE International Conference on **4** (1993), 216 – 219.

[KK06] Noriyuki Kogo and Eiichiro Komatsu, *Angular trispectrum of cmb temperature anisotropy from primordial non-gaussianity with the full radiation transfer function*, Phys. Rev. D **73** (2006), 083007–083012.

[KSH11] M. Kamionkowski, T. L. Smith, and A. Heavens, *Cmb bispectrum, trispectrum, non-gaussianity, and the cramer-rao bound*, Phys. Rev. D **83** (2011), 023007.

[Leo99] N. Leonenko, *Limit theorems for random fields with singular spectrum*, Mathematics and its Applications, vol. 465, Kluwer Academic Publishers, Dordrecht, 1999.

[Lou06] James. D. Louck, *Springer handbook of atomic, molecular, and optical physics, angular momentum theory*, Drake, (Ed.), Springer Science+Business Media, Inc., New York, 2006.

[Mar04] Domenico Marinucci, *Testing for non-Gaussianity on cosmic microwave background radiation: a review.*, Stat. Sci. **19** (2004), no. 2, 294–307 (English).

[Mar06] D. Marinucci, *High-resolution asymptotics for the angular bispectrum of spherical random fields*, Ann. Statist. **34** (2006), 1–41.

[Mar08] Domenico Marinucci, *A central limit theorem and higher order results for the angular bispectrum*, Probability Theory and Related Fields **141** (2008), 389–409, 10.1007/s00440-007-0088-8.

[Mat86] B. Matern, *Spatial variation, lecture notes in statistics, vol. 36*, Springer-Verlag, New York, 2nd ed., 1986.

[McL86] M. G. McLeod, *Stochastic processes on a sphere*, Physics of the Earth and Planetary Interiors **43** (1986), 283299.

[MH95] J. W. D. Molle and M. J. Hinich, *Trispectral analysis of stationary random time series*, Acoustical Society of America **97** (1995), no. 5, 2963–2978.

[Ml66] C. Mller, *Spherical harmonics*, Springer-Verlag, Berlin, Lecture Notes in Mathematics, vol.17, 1966.

[MP10] D. Marinucci and G. Peccati, *Representations of so(3) and angular polyspectra*, Journal of Multivariate Analysis **101** (2010), 77–100.

[MP11] ______, *Random fields on the sphere*, London Mathematical Society Lecture Notes Series, vol. 389, cambridge university press, Cambridge, 2011.

[NC81] G. R. North and R. F. Cahalan, *Predictability in a solvable stochastic climate model*, Journal of the Atmospheric Sciences **38** (1981), 504–513.

[Obu47] A. M. Obukhov, *Statistically homogeneous fields on a sphere*, Usp. Mat. Nauk **2** (1947), 196–198.

[OH02] T. Okamoto and W. Hu, *Angular trispectra of cmb temperature and polarization*, Phys. Rev. D **66** (2002), 063008.

[PBM86] A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev, *Integrals and series. Vol. 2*, Gordon & Breach Science Publishers, New York, 1986, Special functions, Translated from the Russian by N. M. Queen.

[RHS$^+$05] G. Rocha, M. P. Hobson, S. Smith, P. Ferreira, and A. Challinor, *Simulation of non-gaussian cosmic microwave background maps*, Monthly Notices of the Royal Astronomical Society **357** (2005), no. 1, 1–11.

[RST06] S. Rao Jammalamadaka, T. Subba Rao, and Gy. Terdik, *Higher order cumulants of random vectors and applications to statistical inference and time series*, Sankhya (A, Methodology) **68** (2006), 326–356.

[RvN65] M. Rosenblatt and J. W. van Ness, *Estimation of the bispectrum*, Ann. Math. Statist. **36** (1965), 1120–1135.

[Sch42] I. J. Schoenberg, *Positive definite functions on spheres*, Duke Math. J. **9** (1942), 96–108.

[Sch97] M. Schreiner, *On a new condition for strictly positive definite functions on spheres*, Proceedings of the American Mathematical Society **125** (1997), 531–539.

[SG05] O. Schabenberger and C. A. Gotway, *Statistical methods for spatial data analysis*, CRC Press, 2005.

[She11] M. Sherman, *Spatial statistics and spatio-temporal data: Covariance functions and directional properties (wiley series in probability and statistics)*, John Wiley & Sons, Ltd, 2011.

[SK12] T. L. Smith and M. Kamionkowski, *Probability distribution for non-gaussianity estimators constructed from the cmb trispectrum*, Phys. Rev. D **86** (2012), 063009.

[SRG84] T. Subba Rao and M. M. Gabr, *An introduction to bispectral analysis and bilinear time series models*, Springer-Verlag, Berlin Heidelberg, 1984.

[SRT06] T. Subba Rao and Gy. Terdik, *Multivariate non-linear regression with applications*, Dependence in Probability and Statistics (P. Bertail, P. Doukhan, and P. Soulier, eds.), Springer Verlag, New York, 2006, pp. 431–470.

[SW71] E. M. Stein and G. Weiss, *Introduction to Fourier analysis on Euclidean spaces*, Princeton University Press, Princeton, N.J., 1971, Princeton Mathematical Series, No. 32.

[Ter99] Gy. Terdik, *Bilinear stochastic models and related problems of nonlinear time series analysis; a frequency domain approach*, Lecture Notes in Statistics, vol. 142, Springer Verlag, New York, 1999.

[TM98] Gy. Terdik and J. Máth, *A new test of linearity for time series based on its bispectrum*, Journal of Time Series **19** (1998), no. 6, 737–749.

[VMK88] D. A. Varshalovich, A. N. Moskalev, and V. K. Khersonskii, *Quantum theory of angular momentum*, World Scientific Press, 1988.

[WC01] A. Weaver and P. Courtier, *Correlation modelling on the sphere using a generalized diffusion equation*, Quarterly Journal of the Royal Meteorological Society **127** (2001), 1815–1846.

[Wor] P. Wormer, *Angular momentum theory*, Tech. report, Lecture Notes, University of Nijmegen Toernooiveld, 6525 ED Nijmegen, The Netherlands.

[Yad83] M. Ĭ. Yadrenko, *Spectral theory of random fields*, Optimization Software Inc. Publications Division, New York, 1983, Translated from the Russian.

[Yag61] A.M. Yaglom, *Second-order homogeneous random fields*, Proc. Fourth Berkeley Symp. Math. Statist, and Prob., 2, 1961, pp. 593–622.

[Yag87] A. M. Yaglom, *Correlation theory of stationary related random functions*, Springer-Verlag, New York, 1987.